

Departement für Kleintiere, Klinik für Zoo-, Heim- und Wildtiere
der Vetsuisse-Fakultät Universität Zürich

Direktor: Prof. Dr. med. vet. Jean-Michel Hatt

Arbeit unter Leitung von
PD Dr. med. vet. Marcus Clauss

**Reproductive seasonality in captive wild ruminants: implications for
biogeographical adaptation, photoperiodic control, and life history**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Philipp Zerbe

Tierarzt
aus Offenbach a.M., Deutschland

genehmigt auf Antrag von

PD Dr. med. vet. Marcus Clauss, Referent

Zürich 2012

Departement für Kleintiere, Klinik für Zoo-, Heim- und Wildtiere
der Vetsuisse-Fakultät Universität Zürich

Direktor: Prof. Dr. med. vet. Jean-Michel Hatt

Arbeit unter Leitung von
PD Dr. med. vet. Marcus Clauss

**Reproductive seasonality in captive wild ruminants: implications for
biogeographical adaptation, photoperiodic control, and life history**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Philipp Zerbe

Tierarzt
aus Offenbach a.M., Deutschland

genehmigt auf Antrag von

PD Dr. med. vet. Marcus Clauss, Referent

Zürich 2012

Inhaltsverzeichnis:

Zusammenfassung/Abstract	1
Publikation Biological Reviews	2
Abstract	2
Contents	3
Introduction	3
Methods	7
Results	9
(1) Relationships in the basic ruminant dataset	9
(2) Quantifying seasonality	10
(3) Comparison among ruminant groups	10
(4) Factors associated with seasonality	11
(5) Differences between captivity and the wild	12
(6) Seasonality and gestation period	13
Discussion	14
(1) General characteristics of ruminant species	15
(2) Quantifying seasonality	16
(3) Evolving seasonality	17
(4) Causative factors for seasonality: interspecific comparison	18
(5) Environmental correlates of seasonality: intraspecific comparisons	20
(6) Gestation length and reproductive seasonality	23
Conclusions	23
References	24
Supporting information	28
Lebenslauf	
Danksagung	

Zusammenfassung

Reproduktive Saisonalität in Menschenobhut gehaltener Wildwiederkäuer: Rückschlüsse auf biogeographische Adaptation, photoperiodische Kontrollmechanismen und weitere Anpassungen

Zur quantitativen Beschreibung der Reproduktionsmuster wurden Daten von 110 Wildwiederkäuerarten aus Zoos der gemäßigten Zone verwendet (dabei wurde die Anzahl Tage, an denen 80% aller Geburten stattfanden, als Geburtenpeak-Breite [BPB] definiert). Diese Muster wurden mit verschiedenen biologischen Charakteristika verknüpft und mit denen von freilebenden Tieren verglichen.

Der Breitengrad des natürlichen Verbreitungsgebietes korreliert stark mit dem in Menschenobhut beobachteten BPB. Nur 11% der Spezies wechselten ihr reproduktives Muster zwischen Wildnis und Gefangenschaft, wobei für saisonale Spezies die errechnete Tageslichtlänge zum Zeitpunkt der Konzeption für freilebende und in Menschenobhut gehaltene Populationen gleich war.

Reproduktive Saisonalität erklärt zusätzliche Varianzen im Verhältnis von Körpergewicht und Tragzeit, wobei saisonalere Spezies für ihr Körpergewicht eine kürzere Tragzeit aufweisen. Rückschliessend ist festzuhalten, dass Photoperiodik, speziell die absolute Tageslichtlänge, genetisch fixierter Auslöser für die Fortpflanzung ist, und dass die Plastizität der Tragzeit unterstützend auf die erfolgreiche Verbreitung der Wiederkäuer in höheren Breitengraden wirkte.

Key words: Phänologie, Saisonalität, Reproduktion, Tragzeit, Photoperiodik

Abstract

Reproductive seasonality in captive wild ruminants: implications for biogeographical adaptation, photoperiodic control, and life history

A dataset on 110 wild ruminant species kept in captivity in temperate-zone zoos was used to describe their reproductive patterns quantitatively (determining the birth peak breadth [BPB] as the number of days in which 80% of all births occur); then this pattern was linked to various biological characteristics, and compared with free-ranging animals.

Globally, latitude of natural origin highly correlates with BPB observed in captivity, with species being more seasonal originating from higher latitudes. Kept in zoos, 89% of the species retained their reproductive pattern from the wild. For seasonal species, daylength at the time of conception between free-ranging and captive populations was similar.

Reproductive seasonality explains additional variance in the body mass–gestation period relationship, with more seasonal species having shorter gestation periods for their body size. We conclude that photoperiodism, and in particular absolute daylength, are genetically fixed triggers for reproduction that may be malleable to some extent by body condition, and that plasticity in gestation length is an important facilitator that may partly explain the success of ruminant radiation to high latitudes.

Key words: phenology, seasonality, reproduction, gestation, photoperiodism

Reproductive seasonality in captive wild ruminants: implications for biogeographical adaptation, photoperiodic control, and life history

Philipp Zerbe^{1,2}, Marcus Clauss^{1,*}, Daryl Codron^{1,3,4,5}, Laurie Bingaman Lackey⁶, Eberhard Rensch¹, Jürgen W. Streich⁷, Jean-Michel Hatt¹ and Dennis W. H. Müller^{1,8}

¹ *Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland*

² *Section for Small Animal Reproduction, Clinic for Animal Reproduction, University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland*

³ *Florisbad Quaternary Research, National Museum, Bloemfontein, 9300, South Africa*

⁴ *School of Biological and Conservation Sciences, University of KwaZulu-Natal, Private Bag X01, Scottsville 3209, South Africa*

⁵ *Department of Anthropology, University of Colorado at Boulder, Boulder, CO 80309, USA*

⁶ *International Species Information System, 2600 Eagan Woods Drive, Suite 50, Eagan, MN 55121-1170, USA*

⁷ *Leibniz-Institute for Zoo and Wildlife Research, Alfred-Kowalke-Strasse 17, 10315 Berlin, Germany*

⁸ *National Park 'Bavarian Forest', Grafenau, Germany*

ABSTRACT

Many ruminant species show seasonal patterns of reproduction. Causes for this are widely debated, and include adaptations to seasonal availability of resources (with cues either from body condition in more tropical, or from photoperiodism in higher latitude habitats) and/or defence strategies against predators. Conclusions so far are limited to datasets with less than 30 species. Here, we use a dataset on 110 wild ruminant species kept in captivity in temperate-zone zoos to describe their reproductive patterns quantitatively [determining the birth peak breadth (BPB) as the number of days in which 80% of all births occur]; then we link this pattern to various biological characteristics [latitude of origin, mother-young-relationship (hider/follower), proportion of grass in the natural diet (grazer/browser), sexual size dimorphism/mating system], and compare it with reports for free-ranging animals. When comparing taxonomic subgroups, variance in BPB is highly correlated to the minimum, but not the maximum BPB, suggesting that a high BPB (i.e. an aseasonal reproductive pattern) is the plesiomorphic character in ruminants. Globally, latitude of natural origin is highly correlated to the BPB observed in captivity, supporting an overruling impact of photoperiodism on ruminant reproduction. Feeding type has no additional influence; the hider/follower dichotomy, associated with the anti-predator strategy of 'swamping', has additional influence in the subset of African species only. Sexual size dimorphism and mating system are marginally associated with the BPB, potentially indicating a facilitation of polygamy under seasonal conditions. The difference in the calculated Julian date of conception between captive populations and that reported for free-ranging ones corresponds to the one expected if absolute day length was the main trigger in highly seasonal species: calculated day length at the time of conception between free-ranging and captive populations followed a $y = x$ relationship. Only 11 species (all originating from lower latitudes) were considered to change their reproductive pattern distinctively between the wild and captivity, with 10 becoming less seasonal (but not aseasonal) in human care, indicating that seasonality observed in the wild was partly resource-associated. Only one species (*Antidorcas marsupialis*) became more seasonal in captivity, presumably because resource availability in the wild overrules the innate photoperiodic response. Reproductive seasonality explains additional variance in the body mass–gestation period relationship, with more seasonal species having shorter gestation periods for their body size. We conclude that photoperiodism, and in particular absolute day length, are genetically fixed triggers for reproduction that may be malleable to some extent by

* Address for correspondence (E-mail: mclauss@vetclinics.uzh.ch).

body condition, and that plasticity in gestation length is an important facilitator that may partly explain the success of ruminant radiation to high latitudes. Evidence for an anti-predator strategy involving seasonal reproduction is limited to African species. Reproductive seasonality following rainfall patterns may not be an adaptation to give birth in periods of high resource availability but an adaptation to allow conception only at times of good body condition.

Key words: phenology, seasonality, reproduction, gestation, photoperiodism, Bergmann's rule, Rensch's rule.

CONTENTS

I. Introduction	966
II. Methods	970
III. Results	972
(1) Relationships in the basic ruminant dataset	972
(2) Quantifying seasonality	973
(3) Comparison among ruminant groups	973
(4) Factors associated with seasonality	974
(5) Differences between captivity and the wild	975
(6) Seasonality and gestation period	976
IV. Discussion	977
(1) General characteristics of ruminant species	978
(2) Quantifying seasonality	979
(3) Evolving seasonality	980
(4) Causative factors for seasonality: interspecific comparisons	981
(a) Latitude	981
(b) Altitude	982
(c) Predator avoidance	982
(d) Natural diet	983
(e) Mating type	983
(5) Environmental correlates of seasonality: intraspecific comparisons	983
(a) Photoperiodism	983
(b) Resource availability	984
(c) Specific examples	985
(6) Gestation length and reproductive seasonality	986
V. Conclusions	986
VI. Acknowledgements	987
VII. References	987
VIII. Supporting Information	990

I. INTRODUCTION

Habitats do not offer steady conditions to their inhabitants - they are exposed to climatic variations and environmental disasters. Some of these variations are unpredictable incidents such as cosmic or tectonic events, and life reacts to such phenomena rather than adapting to them. Other variations occur more or less regularly, but are nevertheless unpredictable - like the El Niño effect, or fruit mast years. To cope with such variations, species evolve mechanisms that allow for a plastic response of their behavioural patterns. A prerequisite of such behavioural plasticity is a fine detection mechanism that allows early sensing of the emerging event—especially if a response in reproductive output is adaptive. Examples for such adaptations are the reproductive patterns in kakapo (*Strigops habroptilus*) (Powlesland *et al.*, 1992) or dormice (*Glis glis*) (reviewed in Ruf *et al.*, 2006) that only reproduce during fruit mast years. By contrast,

the whole life cycle of a species adapts (and does not just react) to regular and more or less predictable (periodical) environmental variations. One of the most important and best predictable variations is the seasonality in temperate zones, and its most reliable predictor is the variation in day length—photoperiodism—over the year (Bradshaw & Holzapfel, 2007). As an adaptation to seasonal changing of climate conditions within the temperate zone over the year, various animal species of all mammalian clades developed seasonal patterns of reproduction (Bronson, 1989). This allows for offspring to be born when food is available in abundance (which is important to ensure milk production), and the newborns to grow under favourable climatic conditions (Rutberg, 1984; Sinclair, Mduma & Arcese, 2000b; Santiago-Moreno *et al.*, 2006). Detailed studies have demonstrated that in several seasonal wild ruminants, earlier-born offspring can gain more weight and therefore survive the next winter better than later-born offspring (reviewed e.g. by

Côté & Festa-Bianchet (2001)). This seasonal reproduction is cued by photoperiodism (Bradshaw & Holzapfel, 2007), which, *via* the induction of melatonin release, modulates circadian and circannual pacemakers (Hazlerigg & Loudon, 2008; Dardene, 2012). Such physiological pacemakers are an active field of current research (e.g. Lincoln *et al.*, 2006; Hazlerigg & Lincoln, 2011; Hut & Beersma, 2011), but are not the topic of this review.

The seasonal reproduction of certain domestic species has important consequences for animal farming, as the reproduction rate of strictly seasonal species is more or less restricted to one reproductive cycle per year under the natural photoperiodic conditions of the northern hemisphere (Asher, Monfort & Wemmer, 1999; Rhind, Archer & Adam, 2002; Chemineau *et al.*, 2008). To overcome this limitation, artificial photoperiodic treatments are used to control reproduction in seasonal farm animals like sheep, goats and horses (Chemineau *et al.*, 2008). Reproductive seasonality must also be considered in conservation projects of endangered species. A thorough knowledge of the reproductive physiology (including seasonality) is a prerequisite for the successful use of assisted reproduction methods (gamete cryopreservation, artificial insemination, embryo transfer and *in vitro* fertilization) (Jabbour, Hayssen & Bruford, 1997; Sinclair, Ludwig & Clark, 2000a; Penfold, 2005; Santiago-Moreno *et al.*, 2006). The most famous example is the *ex situ* breeding program for the endangered giant panda (*Ailuropoda melanoleuca*), where monitoring of the female oestrus cycle during the short breeding season is used to decide when to introduce the male to the female, or when artificial insemination should be performed (Hodges *et al.*, 1984; Masui *et al.*, 1989).

To detect seasonal reproductive behaviour in the wild, the annual distribution of either mating or birthing events has to be observed. This might be difficult for many species that are either elusive or live in remote habitats, like duikers (Wilson, 2005) or chevrotains (Barrette, 1987; Nowak, 1991; Matsubayashi, Bosi & Kohshima, 2003). Birth records from zoological gardens offer very precise information about reproductive seasonality (Zuckerman, 1952; Brand, 1963; Pelt, 1967; Dittrich, 1970; Kirkwood, Gaskin & Markham, 1987; Skinner, Moss & Skinner, 2002), but it is not certain whether the reproductive seasonality of species in zoos reflects patterns of their conspecifics in the wild (Fairall, 1968). So far, only some exemplary comparisons of seasonality in reproduction between wild and captive populations have been published. It is described that some mammalian species with a fixed breeding season in temperate latitudes reverse their breeding season when transported to the Southern hemisphere, but demonstrate the same breeding season or an irregular pattern when transported to the tropics (e.g. Marshall, 1937; The Duke of Bedford & Marshall, 1942; Zuckerman, 1952; Brand, 1963; Caughley, 1971; Spinage, 1973; Fletcher, 1974; Paré, Barrette & Prescott, 1996; Skinner *et al.*, 2002). As most zoos are located within the temperate zone of the Northern hemisphere with its distinct seasons, but provide food in sufficient amounts and

quality throughout the year, an analysis of reproductive seasonality of wild species in captivity may provide further insights into the regulating mechanisms of reproductive seasonality. For example, Skinner *et al.* (2002) suggested that the change of the sable antelope (*Hippotragus niger*) from a seasonal breeder in the wild to a non-seasonal breeder in captivity could be explained by the continuous feeding regime in the zoo. Pelt (1967) had already presented a similar observation and interpretation due to a comparison of free-ranging and captive blackbuck (*Antelope cervicapra*), and Piening Schuler *et al.* (2009) made similar observations for several gazelle species. On the other hand, the finding that some gazelles that were considered aseasonal breeders, with a seasonal birthing pattern in the wild putatively due to seasonal fluctuations in forage availability, retained their seasonal reproduction in captivity under constant diet provision (Flower, 1932), led to the conclusion that there actually was an underlying photoperiodic response that determined the observed birthing patterns.

The above studies, as well as observations from the wild, mainly provide categories to describe the seasonality of reproductive behaviour (e.g. seasonal *versus* non-seasonal), or describe the monthly distribution of birth events (e.g. Spinage, 1973; Skinner *et al.*, 2002). Even though quantitative, the monthly distribution of births often does not allow for a comparison between different species, because the degree of seasonality is not sufficiently resolved. A species that may give birth to half of the annual offspring in the last week of March and the other half in the first week of April demonstrates the same monthly distribution of births (i.e. 50% in March and in April) as a species that delivers all the annual offspring in March and April but with an even distribution of birthing during the whole months. The first species is, however, notably 'more' seasonal than the second one. Such analyses of the degree of seasonality across different species are needed to test for factors that influence reproductive seasonality. Unfortunately, the generation of such comparative data requires a very close monitoring of whole populations (i.e. a precise documentation of all births within a population), and sufficient data from the wild are therefore mostly not available.

The situation for data of captive-born animals is completely different. The International Species Information System (ISIS) is an institute of the international zoo community that has collected stock data of participating zoos (approximately 750 worldwide) over the last 35 years. These data include the exact dates of birth of almost all individuals born in captivity since 1980. This collection therefore provides data on birth dates in a previously unknown quantity, facilitating a quantitative analysis of reproductive seasonality.

Here, we introduce an expression of reproductive seasonality as a continuous variable, to increase the power of comparative analyses. The following hypotheses then guided our approach.

- (1) The degree of seasonality in wild ruminants depends on a variety of factors:

- (a) Birth seasons become shorter, i.e. animal species become more seasonal, with increasing latitude of their geographical origin, due to a shortening of the season of favourable conditions to raise offspring with increasing latitude (Rutberg, 1987).
 - (b) The mother-young relationship (hider *versus* follower; Lent, 1974) influences the degree of birth seasonality. It was described that 'follower' species (offspring immediately follows mother after birth, and often joins the herd) have consistently shorter birth seasons when compared with 'hider' species (offspring is hidden during the first days of life) (Rutberg, 1987; Sinclair *et al.*, 2000b). This pattern was interpreted as an adaptation to prevent predation. Estes (1976) showed that wildebeest (*Connochaetes taurinus*) calves suffered a lower mortality when born at the peak of the calving season and suggested therefore that birth synchrony was an anti-predator strategy. Similar findings were reported for caribou (*Rangifer tarandus*) (Dauphiné & McClure, 1974; Adams, Singer & Dale, 1995), moose (*Alces alces*) (Testa, Becker & Lee, 2000), wapiti (*Cervus elaphus*) (Smith & Anderson, 1998), roe deer (*Capreolus capreolus*) (Panzacchi *et al.*, 2008) and pronghorn (*Antilocapra americana*) (Gregg *et al.*, 2001). First, the sheer number of young may prevent predators from taking a significant fraction of all newborns when compared to species with non-seasonal reproduction (instantaneous saturation). Second, many adults breeding synchronously may protect their offspring better using group defence in follower species. Third, the presence of many young in a herd may interfere with a predator's ability to pick and pursue a specific target (confusion). Such adaptations are less relevant for hiding species, whose offspring are less vulnerable to predation when compared to offspring that immediately follows their mothers through more open habitats (Rutberg, 1987). Hiders, by contrast, rely on low encounter rates between predators and neonates. In such species, a short birthing season could even be considered disadvantageous, because it would increase the likelihood that roaming predators encounter neonates at a certain period of time, thus promoting adaptations to increase searching activities in this particular time period (Rutberg, 1987). Hiders should therefore benefit from an asynchronous birthing pattern (Ims, 1990a, b). Because such an asynchronous birthing pattern is unlikely to occur at high latitudes, the hider-follower distinction should be particularly relevant for less-temperate species, such as the African ruminant community.
 - (c) The feeding strategy (browser *versus* grazer) influences the degree of birth seasonality. African grazers should have shorter birth seasons compared to browsers, as grasses in rain-dependent regions tend to experience a shorter flush than the deeper-rooted browse plants. This shorter time span in which high-quality food is available for grazers should select for greater birth synchrony (Leuthold & Leuthold, 1975; Skinner *et al.*, 2002). This theory was rejected by Rutberg (1987) for the categorical variables browser *versus* grazer and seasonal *versus* non-seasonal species for a global dataset; Kiltie (1988) also found no support for this hypothesis in another global dataset. A decisive factor in this respect could be that differences between browsers and grazers might be more pronounced in the African biome, due to specific adaptations to the presence of C4 grasses (Codron *et al.*, 2008a; Codron & Clauss, 2010).
 - (d) The mating system (monogamous *versus* polygamous) is associated with the degree of birth seasonality. Polygamous species should be more seasonal than monogamous species. Monogamous males that live in close proximity to their (single) partner throughout the year do not have to cope with the problem of finding the female when it is in oestrus. By contrast, males of polygamous species attempt to copulate with as many females as possible, resulting in higher sexual competition amongst the males. Thus, it might be advantageous if all females are in oestrus at the same time, limiting the time period of high reproductive investment for the males (Lane *et al.*, 2010); in other words, reproductive seasonality might facilitate polygamous mating systems. Because the degree of sexual dimorphism correlates positively with mating system in ruminants (Alexander *et al.*, 1979; Jarman, 1983; Weckerly, 1998; Loison *et al.*, 1999), we therefore also expect more seasonal species to have a higher degree of male-biased sexual dimorphism.
- (2) Differences between the wild and captivity can help to explain parts of the mechanisms that lead to seasonal reproductive patterns:
- (a) Species that originate from a different hemisphere than the one they are kept in switch their seasonal birth pattern accordingly (Marshall, 1937; Zuckerman, 1952; Brand, 1963; Caughley, 1971; Spinage, 1973; Fletcher, 1974; Lincoln, 1985; Paré *et al.*, 1996; Skinner *et al.*, 2002).
 - (b) Species with a distinct seasonal reproductive pattern, that originate from a different latitude than the one they are kept in, should conceive

when the absolute daylength at the location where they are kept is similar to the absolute day length of breeding in their natural habitat. Because day length changes across the year with latitude, this implies a shift in breeding events in relation to the calendar year (Fig. 1). The direction of this shift depends on two factors—whether the species is kept at lower or higher latitudes than their natural origin, and whether breeding is usually triggered before or after equinox (Fig. 1). Rutberg (1987) predicted such a shift in the timing of reproductive activity between populations at different latitudes. If day length is the triggering photosignal for the initiation of mating or conception, then differences in conception patterns between the wild and captivity should be of such a magnitude that the absolute day length in the wild (x) and in captivity (y) at the initiation of reproductive activity is identical ($y = x$). Evidence for or against such a mechanism so far stems mainly from intraspecific comparisons and is somewhat ambiguous. On the one hand, several authors suggest that within a species, latitude does not influence this pattern (Fletcher, 1974; Paré *et al.*, 1996; Moe, Rutina & Du Toit, 2007); on the other hand, many other authors demonstrate that such a pattern exists (Ransom, 1966; Fairall, 1968; Sadleir, 1969; Nievergelt, 1974; Bunnell, 1982; fig. 5 in Bronson, 1985; Rutberg, 1987 cites several comparisons of populations in East Africa and South Africa; Saether *et al.*, 1996; Flood & Tedesco, 1997; Linnell & Andersen, 1998; Bonenfant *et al.*, 2004; Loe *et al.*, 2005; several examples in Santiago-Moreno *et al.*, 2006; Ungerfeld *et al.*, 2008). In the case of the red deer (*Cervus elaphus*), the direct conflict of reports (Fletcher, 1974; Bonenfant *et al.*, 2004; Loe *et al.*, 2005) makes this question particularly interesting. In the case of the Himalayan tahr (*Hemitragus jemlahicus*), a change of the birthing pattern with latitude would be obvious if one single outlier (out of seven populations) was excluded from the published analysis (*cf.* fig. 2 in Paré *et al.*, 1996). Alternatively, if absolute day length was not important in ruminants, and breeding was mainly triggered by changes in day length only (as commonly assumed for long-lived animals; Bradshaw & Holzapfel, 2007), then no systematic offset of reproductive events between free-ranging and captive populations should be discernible.

- (c) Species expand their breeding season or change from a seasonal to a non-seasonal breeding pattern due to the reduced selective pressure (no predation) or the unlimited water and/or food supply in captivity (Caughley, 1971;




Fig. 1. Day length (including civil twilight) of each day of the year at different latitudes (calculated according to Forsythe *et al.*, 1995). The window of the mating season (boxed area) is indicated for short-day breeders (i.e. animals that mate during the period of the year when day length decreases). If mating was triggered by a day length threshold, we would expect species originating from higher latitudes than the one they are kept at to start reproducing sooner in captivity than in the wild if the day length threshold is before equinox, but later than in captivity if the day length threshold is after equinox. For species originating from a lower latitude than the one they are kept at, the opposite pattern should apply.

Skinner *et al.*, 2002; Piening Schuler *et al.*, 2009). Intraspecific comparisons of birthing patterns between habitats of different quality, between animals of different body condition, or differences in resource availability between years suggest that a higher resource availability, *via* a quicker attainment of a high nutritional status, facilitates earlier mating and birthing seasons, even if the general onset of reproductive activity is still determined by photoperiod (Verme, 1965, 1969; Nievergelt, 1966; Mitchell & Lincoln, 1973; McGinnes & Downing, 1977; Hamilton & Blaxter, 1980; Flydal & Reimers, 2002; Post, 2003). Correlations of rainfall patterns and food quality with mating/birthing events in less pronounced seasonal species support this concept (Moe *et al.*, 2007). Correspondingly, we additionally expect that those species that maintain a seasonal reproductive pattern in captivity enter into reproductive activity somewhat earlier than in the wild due to their putatively better nutritional status; in other words, for the model suggested above with day length in the wild (x) and in captivity (y) at the initiation of reproductive activity, we predict a relationship of $y = 1.0x + a$ due to a systematic forward shift of reproduction due to unlimited food supply. In other species, which show a moderate seasonality in the wild, we expect the high levels of nutrition in captivity to result in disappearance of a seasonal reproduction pattern (Montgomery, Scott & Hudson, 1985).

- (3) Reproductive seasonality is associated with other life-history parameters that help facilitate the respective patterns: seasonal species adjust their gestation period length so that offspring are born at times that ensure maximum offspring survival; therefore, seasonality can partly explain the residual variation in the relationship between body mass and length of the gestation period (Müller *et al.*, 2011*b*). The gestation period of artiodactyls scales to body mass^{0.16±0.03SD} (Western, 1979). The variation in this relationship might derive in part from the fact that larger animals shorten, and smaller animals lengthen their gestation period in relation to body mass in order to achieve birthing at a beneficial time of the year at high latitudes. This concept was elaborated in detail by Kiltie (1984, 1988) who also suggested that species in which gestation times exceed the time between two seasonal periods of high resource availability might be excluded from reproducing (and hence living) competitively at higher latitudes. As this author notes, the (historical) occurrence of species with gestation periods longer than 1 year—perissodactyls, proboscids and camelids—at higher latitudes indicates that a reproductive biology characterised by short gestation periods is not a *conditio sine qua non*, but should be considered one of many possible adaptations to life at high latitudes.

II. METHODS

We used data of approximately 200 000 animals representing 110 different species of captive ruminants (suborder Ruminantia) from the International Species Information System (ISIS) data files. The observation interval was between 1980 and 2006. We removed species that had their last recorded birth before 1997 (e.g. that were no longer kept in ISIS zoos), or had an overall birth-count below 90. Only records with an exact birth date and from countries of the northern hemisphere with distinct seasons (spring, summer, autumn and winter) were used, i.e. we generally excluded animals from zoological collections of the Arabian Peninsula, Australia, Asia (except for Japan), Mexico, South America and Africa. Exceptions to this rule are explicitly mentioned. Years were sub-divided into 73 consecutive 5-day intervals, and the total number of births that occurred within each interval was counted. For leap years, births from December 31st were allocated to the 73rd interval. As there was a salient increase of births on January 1st, which was most likely the effect of the reporting management of some zoos, the first interval of the year was calculated as the average of the second and preceding 73rd interval. This effect was noticeable in 33 of the 110 species. We are aware of the fact that some zoos proactively time breeding of their animals to prevent births in certain seasons of the year; however, to our knowledge, this strategy is not applied universally, neither on a species nor on a zoo level. On the other hand, zoos may actively prevent breeding by separation or medical

contraception. Although this is rare in wild ruminants, such procedures will impact the reproductive output; however, this should not affect results on reproductive seasonality.

For the comparison of seasonality patterns between the wild and zoo data, we had to introduce a categorical variable in order to accommodate literature data for free-ranging populations. Species were allocated subjectively, based on their birthing patterns, to five categories (Fig. 2): Category 1: narrow peak (less than approximately 60 days), no births for the rest of the year. Category 2: expanded peak (<60 days) or double peak, no births for the rest of the year. Category 3: peak(s) with small number of births throughout the year. Category 4: births throughout the year, undulating pattern (possibly indicating preferred seasons). Category 5: constant births throughout the year without a clear preferred season.

In order to allocate species from captivity to category 4 or 5, at least 150 birth events were required. Due to the concentration of birth events within a rather short time period in more seasonal species (categories 1–3), 90 birth events were considered sufficient for an allocation into these categories (*cf.* a similar observation on the number of birth records required for seasonal and aseasonal species by Zuckerman, 1952).

Data on categorical birth seasonality in the wild, mid-latitude of geographical origin, body mass, sexual size dimorphism, mating type, mother-young relationship, proportion of grass in the natural diet, and the length of the gestation period, were taken from the literature (online Appendix S1). We used data on the start of the birthing period in combination with the gestation length of the respective species to calculate the start of the mating/conception season in captivity, and in the wild. Because transitions between the seasonality categories are to a degree subjective, a change of more than 1 category was considered as a true change in the seasonality pattern when comparing seasonality categories between free-ranging and captive populations. The day length of a specific day of the year, at a specific latitude, was calculated, including civil twilight (which is defined as the time between sunrise or sunset and when the centre of the sun is six degrees below the horizon; light during civil twilight is bright enough to perform ordinary outdoor activities without artificial light), according to Forsythe *et al.* (1995). For these calculations, the mean latitude of captive populations was assumed to be 45°. For the overall relationship with reproductive seasonality, we used the median latitude of origin for species given in the PanTHERIA database (Jones *et al.*, 2009; online Appendix S1); we appreciate that using the median of the range given in this database introduces a source of error in the dataset, because the core area of origin for many species will not be in the middle of their reported range; it should be noted, however, that this source of error is not *a priori* systematic. For the comparison of the day length at which mating/conception occurred in the wild and in captivity (which included only species with a BPB80 < 33 days; see Section III.5), we used data on individual free-ranging populations from the literature for which the exact latitude was noted (online Appendix S2). The date of mating/conception was calculated using literature




Fig. 2. Classification of seasonality categories according to the pattern of birth distributions across the year (as divided into 5-day blocks) from 1: highly seasonal to 5: aseasonal species. 1, *Dama dama*; 2, *Connochaetes taurinus*; 3, *Nanger soemmeringii*; 4, *Nanger dama*; 5, *Tragulus javanicus*.

reports on the birthing pattern of the species and re-calculating conception by subtracting the gestation period. The literature sources for Appendices S1–3 are collated in Appendix S4.

As a quantitative measure of the degree of reproductive seasonality, we defined the birth peak breadth (BPB) as the number of days (calculated as 5-day intervals) in which a certain percentage (50, 70, 80 and 90%; BPB50, BPB70, BPB80 and BPB90, respectively) of all births occurred (adapted from Rutberg, 1987). A command line script ('peakfinder', cross platform Perl script) was developed to search for the smallest number of successive intervals that included the demanded percentage of births (Fig. 3). When there was more than one such birth peak (i.e. same number of 5-day intervals), we used the window with the highest number of births for further analyses.

Comparisons between ruminant subfamilies were performed using a one-way analysis of variance (ANOVA) with *post hoc* tests using Sidak adjustment for multiple comparisons. Nonparametric correlations were analyzed by Spearman's ρ and parametric correlations by Pearson's R . To analyze the influence of external and physiological factors on seasonality, body mass, latitude of geographical origin, mating type or sexual dimorphism, mother–young relationship, proportion of grass in the natural diet, and whether or not a species was of African origin, we included and then excluded these factors or covariates in a backward stepwise general linear model approach, including two-way interactions. Because the interaction of African species and latitude, and the interaction of African species and the mother–young relationship were significant in the resulting model (data not shown), and because several predictions




Fig. 3. Different approaches to determine the degree of reproductive seasonality in Père David's deer (*Elaphurus davidianus*). The dotted lines indicate start and end of the 80% birth approach by Rutberg (1987), defined as the time period in which 80% of all births take place from the beginning of the first birthing activity onwards, and the solid gray lines delineate the shaded gray area that denotes the birth peak breadth (BPB80), defined as the shortest time period in which 80% of all births take place. Note the better representation of the real reproductive pattern by the BPB approach.

referred to African or non-African species specifically, we decided to perform the following analyses on three datasets—the total dataset, African species, and non-African species. We report results of various linear models in which specific cofactors were tested; we also report the final linear model for every dataset in which non-significant factors were successively excluded (without interactions). To achieve normality, body mass was log-transformed in advance. To test the hypothesis that animals evolved changes in their gestation period length to adapt to a seasonal environment, another backward stepwise linear model was used in which the length of the gestation period was the dependent variable, and body mass, seasonality (as BPB80), mating type or sexual dimorphism, mother-young relationship, and proportion of grass in the natural diet were included as independent variables or cofactors, respectively. We used the relative gestation length (in days body mass^{-0.12}; exponent derived from linear regression analysis from log-transformed gestation length and body mass) for a graphical representation of the results.

In order to avoid false interpretation of ancestry-based correlations in these models as adaptations (i.e. finding a significant result simply because related species behave in a similar manner) (Felsenstein, 1985; Pagel, 1999), the analyses were not only performed as ordinary least squares (OLS), but also controlled for phylogenetic influences using the phylogenetic generalized least-squares method (PGLS) (Martins & Hansen, 1997; Rohlf, 2001). The phylogenetic tree was derived by pruning the mammal supertree from Bininda-Emonds *et al.* (2007) for those species not represented in the dataset, using Mesquite (Maddison & Maddison, 2006). Missing taxa were included [*Gazella arabica*] was managed as one population by the zoo community but is in fact a hybrid population; *Ovis orientalis* was split into

three subspecies (*O.o.musimon*, *O.o.gmelini*, *O.o.vignei*) as the start of their breeding seasons differ markedly from each other] and polytomies were resolved (see online Appendix S3 for the phylogenetic tree used). Because the resulting tree was not based on our own calculations of branch lengths using consistently the same characters, we used only a tree without branch lengths. The statistical calculations were performed with PASW 18.0 (SPSS Inc., Chicago, IL, USA) and COMPARE 4.6 program (Martins, 2004). The significance level was set to $P < 0.05$. Differences in significance between OLS and PGLS are indicated in results tables by grey shading.

III. RESULTS

(1) Relationships in the basic ruminant dataset

In our dataset, there was no significant relationship between (log-transformed) body mass and latitude of natural origin across species in the whole dataset or among African species (Table 1). For non-African species, the relationship was not significant using OLS, but was significant when phylogenetic relationships were accounted for (Table 1); in other words, although the overall relationship was not significant when assuming that all datapoints are independent (OLS), there is an increase in body mass with increasing latitude within clusters of more closely related species (among non-African ruminants)—which corresponds to the predictions made by Bergmann's rule (see Section IV.1).

Sexual size dimorphism increased with increasing body mass in the overall dataset and among non-African species (Table 1), corresponding to Rensch's rule (see Section IV.1). By contrast, the relationship was not significant using PGLS in African species (Table 1), indicating that the relationship does not occur equally in different taxonomic groups. Body mass was positively related to the percentage of grass in the natural diet in the overall dataset and among non-African species, but not among African species when accounting for phylogeny (Table 1). Accounting for phylogeny, an increase of sexual size dimorphism tended ($P = 0.053$) to occur with increasing latitude in the overall dataset but not in the African and non-African subsets (Table 1). Latitude was not correlated to the percentage of grass in the natural diet, but correlated negatively with the length of the gestation period in the overall dataset and among African species in OLS; using PGLS, there was a corresponding trend ($P = 0.056$) in the overall dataset (Table 1). The percentage of grass in the natural diet increased with increasing sexual size dimorphism in the overall dataset and among non-African species, but not among African species (Table 1).

The allometric equations for the length of the gestation period (Gestlength), determined using log-transformed data, were, with 95% confidence intervals in parentheses,

OLS: Gestlength (days) = 131 (117–146) $BM^{0.12(0.09-0.14)}$ ($R = 0.67$, $P < 0.001$)

PGLS: Gestlength (days) = 145 (120–174) $BM^{0.10(0.08-0.12)}$ ($R = 0.61$, $P < 0.001$).

Table 1. Correlations between the continuous measurements in the basic ruminant dataset used in this study (data from online Appendix S1)

Correlation	Model	All species ($N = 110$)		African species ($N = 47$)		Non-African species ($N = 63$)	
		R	P	R	P	R	P
BM—Lat	OLS	0.07	0.444	−0.08	0.603	0.16	0.203
	PGLS	0.12	0.213	−0.14	0.336	0.27	0.032
BM - BM m/f	OLS	0.42	<0.001	0.33	0.024	0.50	<0.001
	PGLS	0.36	<0.001	0.12	0.407	0.45	<0.001
BM - %grass	OLS	0.38	<0.001	0.29	0.047	0.47	<0.001
	PGLS	0.20	0.039	0.21	0.157	0.27	0.032
Lat - BM m/f	OLS	0.27	0.004	0.27	0.070	0.17	0.185
	PGLS	0.19	0.053	0.24	0.100	0.21	0.103
Lat - %grass	OLS	0.09	0.334	0.27	0.066	0.06	0.647
	PGLS	0.05	0.635	0.22	0.145	0.01	0.912
Lat - Gestlength	OLS	−0.25	0.007	−0.33	0.024	−0.14	0.273
	PGLS	−0.18	0.056	−0.27	0.065	−0.01	0.912
%grass - BM m/f	OLS	0.34	<0.001	0.13	0.383	0.56	<0.001
	PGLS	0.35	<0.001	0.15	0.312	0.49	<0.001

%grass, percentage of grass in the natural diet; BM m/f, body mass ratio between males and females (a measure for sexual size dimorphism); BM, body mass (log-transformed); Gestlength, length of gestation period; Lat, latitude of natural origin; OLS, ordinary least squares; PGLS, phylogenetic generalized least squares (accounting for phylogenetic relationships between species; differences to OLS indicated by grey shading).

(2) Quantifying seasonality

The correlation of the ordinal birth season categories with the continuous variables BPB50, 70, 80 and 90 in our data was significant ($P < 0.001$ in all cases), indicating that the quantitative approach of the BPB and the categorical approach necessary to classify literature data yields comparable results. All BPB variables showed a bimodal distribution across species; the larger the BPB chosen (from 50 to 90), the more this distribution was shifted to the right (Fig. 4). This bimodality suggests that although intermediate forms of reproductive seasonality exist, both strict seasonality and distinct aseasonality are the most common modes in ruminants. The smallest BPB50 was 10 days, the smallest BPB70 20 days, the smallest BPB80 30 days, and the smallest BPB90 40 days. Whereas the beginning of the birthing season in captivity did not show a systematic pattern in species originating from lower latitudes, the beginning of birthing seasons was similar for species originating from higher latitudes (Fig. 5).

(3) Comparison among ruminant groups

There were significant differences both in the mean latitude of origin and the degree of seasonality among ruminant taxonomic groups (Table 2). In particular, the extant Capreolinae (New World cervids) and the Caprinae originate from higher latitudes than many other ruminant taxa, and also have shorter BPBs. When correlating the minimum and maximum latitude of origin and BPB80 with the variance of the respective measurements for the taxonomic groups, a distinct pattern is evident: whereas the minimum latitude and the maximum BPB80 were not correlated with the variance of their respective measurements




Fig. 4. Distribution of species according to their reproductive seasonality, characterised by the birth peak breadth (BPB, in days) in which (A) 50%, (B) 70%, (C) 80%, (D) 90% of all births occur.

(latitude: $r = 0.33$, $P = 0.290$; BPB80: $r = 0.10$, $P = 0.760$), the maximum latitude and the minimum BPB80 were both highly correlated (latitude: $r = 0.89$, $P < 0.001$; BPB80: $r = -0.85$, $P = 0.001$; Fig. 6), indicating that low-latitude origin and a low degree of reproductive seasonality did not




Fig. 5. Relationship between the (absolute) median latitude of origin of 110 ruminant species and the range of Julian days during which 80% of all births occur in captivity (BPB80). Note that for some more tropical species BPB80 reaches from one year to the next (crossing Julian day 365). For species from higher latitudes, the time of the birth window has a comparatively consistent start, and the birth window is shorter.

increase the overall range of these parameters covered by a taxonomic group, but that it was the inclusion of higher latitude origins and more distinct reproductive seasonality that increases the overall range of latitudes and seasonality present in a taxonomic group.

(4) Factors associated with seasonality

In the following, results for the BPB80 are given. Results for other BPBs are only reported if they deviate from those of BPB80.

The BPB80 was highly related to the latitude of geographic origin in the overall dataset, and among African and non-African species (Table 3). When plotting species on the world map according to their seasonality category, the relationship with latitude is evident (Fig. 7). Body mass was

not correlated to any BPB (data not shown) and did not contribute significantly to the relationship between latitude and BPB80 (Table 4).

With respect to an influence of the mother-young-relationship on the effect of latitude on the BPB, there was a remarkable difference between the overall dataset and the non-African species on the one hand, with no additional effect of the hider/follower dichotomy, and the African species on the other hand (Table 5). In the African species, the hider/follower dichotomy had a significant effect on the BPB80, and in the PGLS model the inclusion of the hider/follower dichotomy reduced the effect of latitude on BPB80 to non-significance (Table 5).

The percentage of grass in the natural diet had no influence on the relationship between latitude and BPB80, neither in the overall dataset, nor among the African or non-African species, respectively (Table 6).

The effect of the mating system was tested in two different ways—using the mating-system categories ‘monogamous’, ‘tending’ and ‘harem’ as cofactors, or using the ratio of male to female body mass, which indicates the sexual size dimorphism, as a continuous covariable. For BPB80, mating system did not have a significant effect (data not shown), but for BPB50, an additional effect of mating system was significant in the overall dataset (Fig. 8); this significance was more pronounced with PGLS, indicating that the effect occurred in several ruminant lineages (Table 7). Sexual size dimorphism tended to have an effect on the relationship of latitude and the BPB80 in all datasets (Table 7).

For the overall dataset and the non-African species, the final model after backwards removal of non-significant factors resulted in only latitude having a significant effect on BPB80 (Table 3); for the African species, this procedure resulted in the model that included latitude and mother-young-relationship (Table 5).

Table 2. Mean \pm standard deviation (range in parentheses) of the latitude of origin and the birth peak breadth in which 80% of all births occur (BPB80, in days) in wild ruminant species sorted according to taxonomic groups

Taxon	N	Latitude	BPB80
Tragulidae	2	6 (4–7) ^{ab}	285 (285–285) ^{ab}
Antilocapridae	1	37	45
Giraffidae	2	3 (2–5) ^a	265 (255–275) ^{ab}
Moschidae	1	53	50
Capreolinae	9	38 \pm 19 (10–65) ^{bc}	90 \pm 80 (35–280) ^c
Cervinae	13	26 \pm 13 (7–48) ^{ac}	145 \pm 85 (45–270) ^{ac}
Muntiacinae	3	22 \pm 9 (12–28) ^{ac}	250 \pm 40 (205–280) ^{ab}
Cephalophinae	4	6 \pm 6 (1–12) ^a	265 \pm 10 (255–275) ^b
Bovinae	18	16 \pm 14 (1–51) ^a	230 \pm 50 (115–290) ^b
Reduncinae	5	9 \pm 4 (6–14) ^a	255 \pm 30 (205–280) ^{ab}
Aepycerotinae	1	13	190
Antilopinae	18	16 \pm 12 (0–46) ^a	230 \pm 70 (30–275) ^b
Hippotraginae	6	14 \pm 7 (5–21) ^a	255 \pm 15 (240–270) ^b
Alcelaphinae	5	19 \pm 10 (6–30) ^{ac}	150 \pm 35 (110–195) ^{bc}
Caprinae	22	39 \pm 13 (10–72) ^c	80 \pm 55 (30–260) ^c

Means with different superscripts within a column differ significantly (ANOVA, Sidak *post hoc* test).




Fig. 6. Relationship between the (A) minimum and (B) maximum latitude of origin and the (C) minimum and (D) maximum of the birth peak breadth in which 80% of all births occurred (BPB80) with the variance of the respective measurements (data from Table 2) for different ruminant taxonomic groups. Note the clear pattern that among taxonomic groups, the variance in latitude increases with maximum latitude, and the variance in BPB80 is higher for lower minimum BPB80, suggesting that taxonomic groups expanded into higher latitudes, and evolved shorter BPB80, from lower latitudes (with longer BPB80).

Table 3. Correlations of the birth peak breadth in which 80% of all births occur (in days) and latitude of geographic origin in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using ordinary least squares (OLS) and phylogenetic generalized least squares (PGLS)

Dataset	Model	R	P
All species	OLS	-0.82	<0.001
	PGLS	-0.78	<0.001
African species	OLS	-0.52	<0.001
	PGLS	-0.34	0.020
Non-African species	OLS	-0.79	<0.001
	PGLS	-0.75	<0.001

(5) Differences between captivity and the wild

Eleven out of 103 species (10.7%) changed their pattern of reproductive seasonality between the wild and captivity by more than one category. These were (category free ranging/captive) *Axis porcinus* (2/4), *Cervus alfredi* (2/4), *Cervus unicolor* (2/4), *Bos javanicus* (2/4), *Bubalus bubalis* (2/4), *Kobus megaceros* (2/4), *Antidorcas marsupialis* (4/2), *Antilope cervicapra* (2/4), *Gazella dorcas* (2/4), *Hippotragus niger* (2/4), and *Hemitragus hylocrius* (2/5). Ten of these changed to a less seasonal pattern in captivity; only the springbok (*Antidorcas marsupialis*) had a more seasonal pattern in zoos.

All these species had their median origin at latitudes $\leq 23^\circ$, indicating that substantial changes in the seasonal pattern of reproduction occurred in low-latitude species only.

The only two seasonal species from the southern hemisphere for which exact data from the wild are available - *Damaliscus lunatus* and *Connochaetes gnou* - shifted their birth season start by nearly half a year (start of BPB80: 155 days earlier than data from the wild in both cases).

Comparing the beginning of the mating/conception season in free-ranging and captive populations of clearly seasonal (BPB80 < 33 days) species (data for captive species from online Appendix S1, for free-ranging populations from online Appendix S2; $N = 28$), there was a significant correlation between day length on the day reproductive activity was initiated in the wild and in captivity (Fig. 9). The slope of the linear regression line included 1.0 in the 95% confidence interval for the whole dataset in the PGLS analysis, and after excluding five visible outliers in both the OLS and the PGLS analysis (Table 8), indicating that the day lengths at the beginning of reproductive activity in the wild and in captivity were mostly identical, and that the shifts in the start of reproductive activity exemplified in Fig. 1 actually occurred (Fig. 10). The numerically positive intercept of the regression equation was not significantly different from zero (Table 8). Among the species considered outliers (see Fig. 9)



Fig. 7. Distribution of ruminant species across the world. Location of dots indicates the median of the natural origin; grey shading codes categorical seasonality. The degree of grey shading of dots (from white to black) correlates negatively with the length of the birth peak breadth in which 80% of all births occur (BPB80, in days), i.e. aseasonal species are represented by white dots (BPB80 = 240–290 days), strictly seasonal species by black dots (BPB80 = 30–80 days), with three intermediate shading steps (covering the BPB80 ranges of 180–235 days, 135–175 days and 85–125 days, respectively).

Table 4. Results from linear models testing for an additional effect of (log-transformed) body mass on the relationship between latitude of geographical origin and the birth peak breadth in which 80% of all births occur (in days) in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using ordinary least squares (OLS) and phylogenetic generalized least squares (PGLS)

Dataset	Model	R^2	Latitude		Body mass	
			F/t	P	F/t	P
All species	OLS	0.68	222.248	<0.001	0.199	0.656
	PGLS	0.55	11.143	<0.001	0.227	0.821
African species	OLS	0.31	17.749	<0.001	2.642	0.111
	PGLS	0.12	2.500	0.016	0.300	0.765
Non-African species	OLS	0.63	99.073	<0.001	0.525	0.472
	PGLS	0.57	8.700	<0.001	0.704	0.484

were the muskoxen (*Ovibos moschatus*), the bighorn sheep (*Ovis canadensis*) and the Dall sheep (*Ovis dalli*), whose populations start breeding at day lengths in the wild (19.9, 8.7 and 8.4 h, respectively) that are never achieved at the average latitude of the zoological institutions in our dataset (45° north). When kept at a higher latitude (than the origin), the onset of the mating season is delayed in some species, leading to later births, and when kept at lower latitude (than the origin), the onset of the mating season is earlier in some species, leading to earlier births. In the goitered gazelle (*Gazella subgutturosa*), for which data were available from Arabian and European zoological facilities, a corresponding shift in the birthing season was evident between the two captive populations (Fig. 11).

Table 5. Results from linear models testing for an additional effect of the mother-young-relationship (hider/follower) on the relationship between latitude of geographical origin and the birth peak breadth in which 80% of all births occur (in days) in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using ordinary least squares (OLS) and phylogenetic generalized least squares (PGLS; differences to OLS indicated by grey shading)

Dataset	Model	R^2	Latitude		Hider/follower	
			F/t	P	F/t	P
All species	OLS	0.68	157.416	<0.001	2.072	0.153
	PGLS	0.56	10.857	<0.001	0.968	0.335
African species	OLS	0.52	4.129	0.048	23.296	<0.001
	PGLS	0.42	1.733	0.090	4.049	<0.001
Non-African species	OLS	0.62	88.229	<0.001	0.029	0.865
	PGLS	0.75	8.600	<0.001	0.235	0.815

(6) Seasonality and gestation period

The relative gestation period showed a positive correlation with BPB in both OLS and PGLS, indicating that more seasonal species had shorter gestation periods (BPB80: OLS $\rho = 0.49$, $P < 0.001$; PGLS: $r = 0.36$, $P < 0.001$). For the absolute length of the gestation period, body mass, BPB80, the mother-young-relationship and sexual dimorphism (but not whether species were of African or non-African origin) were significant influence factors in OLS in the whole dataset. In PGLS, only body mass, BPB80 and sexual size dimorphism were significant (Table 9). For African species, body mass and BPB80 were the only significant covariables in PGLS (Table 9). Among non-African species, the inclusion or exclusion of the roe deer (*Capreolus capreolus*)—known to

Table 6. Results from linear models testing for an additional effect of the percentage of grass in the natural diet on the relationship between latitude of geographical origin and the birth peak breadth in which 80% of all births occur (in days) in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using ordinary least squares (OLS) and phylogenetic generalized least squares (PGLS)

Dataset	Model	R^2	Latitude		%grass	
			F/t	P	F/t	P
All species	OLS	0.68	221.582	<0.001	1.244	0.267
	PGLS	0.55	11.000	<0.001	1.250	0.214
African species	OLS	0.28	12.907	0.001	1.146	0.290
	PGLS	0.12	2.500	0.016	0.500	0.620
Non-African species	OLS	0.63	98.737	<0.001	1.101	0.298
	PGLS	0.76	8.600	<0.001	1.000	0.321

have delayed implantation and therefore a particularly long gestation period—had a distinct effect on the results of the analyses (Table 9). Without the roe deer, BPB80 was the only significant factor influencing the gestation period in addition to body mass in PGLS; when the roe deer was included, however, the influence of BPB80 was not significant in PGLS, whereas now that of the mother-young-relationship as well as that of sexual dimorphism was. Generally, the interaction between body mass and BPB80 was not significant (not shown), indicating no evidence for a change of the influence of BPB80 across the body size range. Ruminants with a more distinct seasonal reproduction had, on average, shorter gestation periods than similar-sized less-seasonal ruminants (Fig. 12A); in other words, species with a more pronounced seasonal reproduction pattern often had relatively short gestation periods for their body mass. This pattern was evident in several ruminant lineages (Fig. 12B, C). There were some notable outliers to this pattern. Giraffids have longer gestation periods than other ruminants (Fig. 12A, B; Müller *et al.*, 2011b). The roe deer (*Capreolus capreolus*) had a surprisingly long gestation period for a seasonal (and for any) ruminant of its body size. The pronghorn (*Antilocapra americana*), the pudu (*Pudu pudu*), and the Père David's deer (*Elaphurus davidianus*) also had relatively long gestation periods for ruminants of their body size range with a distinct seasonal reproduction pattern (Fig. 12A, B).

IV. DISCUSSION

This study demonstrates that in contrast to warning against the use of data from captive populations for the characterisation of the seasonality of reproductive events—because they differ from those in the wild (Fairall, 1968), such data can be used fruitfully, partly even *because* of a systematic difference to events in free-ranging populations. The degree of reproductive seasonality of a species can be characterised by a continuous parameter, the birth peak breadth (BPB, the number




Fig. 8. Relationship between birth peak breadth in which 50% of all births occur in captivity (BPB50, in days) and the latitude of a species' origin. Note that species from higher latitudes demonstrate a higher degree of reproductive seasonality when compared to species that originated from lower latitudes.

of days in which a certain percentage of all births occur). This parameter correlates well with the more traditional categorical description of seasonality (highly seasonal, seasonal, aseasonal) and facilitates testing the influence of biological and environmental factors on the degree of seasonality. In particular, we could demonstrate that seasonality of reproduction in ruminants is independent of body mass and mainly linked to the latitude of the origin of the species; that there is a difference between African and non-African species with respect to birth synchrony in the lider/follower strategy; that the photoperiodic signal that many seasonal species adapt to is probably the absolute day length; and that a shortening of the gestation period may be an important adaptation of many

Table 7. Results from linear models testing for an additional effect of the mating system (monogamous/tending/harem) or the sexual size dimorphism on the relationship between latitude of geographical origin and the birth peak breadth (in days) in which 50% (mating system) or 80% (dimorphism) of all births occurred in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using ordinary least squares (OLS) and phylogenetic generalized least squares (PGLS; differences to OLS indicated by grey shading)

Dataset	Model	R^2	Latitude		Mating system	
			F/t	P	F/t	P
All species	OLS	0.65	173.633	<0.001	3.156	0.047
	PGLS	0.55	11.750	<0.001	4.140	0.019
African species	OLS	0.30	11.429	0.002	1.465	0.242
	PGLS	0.18	2.083	0.043	1.800	0.177
Non-African species	OLS	0.64	93.096	<0.001	1.874	0.163
	PGLS	0.78	10.000	<0.001	2.180	0.122
Dimorphism						
All species	OLS	0.70	199.876	<0.001	6.456	0.012
	PGLS	0.58	11.000	<0.001	1.886	0.062
African species	OLS	0.32	12.214	0.001	3.251	0.078
	PGLS	0.19	1.929	0.060	2.054	0.046
Non-African species	OLS	0.64	95.263	<0.001	3.874	0.054
	PGLS	0.60	8.300	<0.001	1.748	0.086




Fig. 9. Relationship between the day length at the beginning of mating/conception in the wild and in captivity (data from online Appendices S1 and S2) in 28 ruminant species. When omitting five outlier species (indicated by open symbols) from the analysis, a linear relationship results (solid line) that is not significantly different from the $y = x$ line (broken line) (for statistics, see Table 8).

seasonal species to ensure birthing at times of favourable conditions but facilitating a rutting season to take place in autumn or even early winter at the same time. In particular, this study ($N = 110$) increases the species number included in comparative analyses on the relevance of seasonal reproduction patterns significantly (Rutberg, 1987: $N = 27$; Kiltie, 1988: $N = 28$ ruminant species), and thus allows a more rigorous testing of additional effects, such as mother-young relationship, diet, or mating type. Our basic data compilation also demonstrated some additional relationships between life-history and ecological variables among ruminants.

(1) General characteristics of ruminant species

Bergmann’s rule states that in individuals of a species, or of closely related species, body mass increases with increasing latitude. Previous analyses of ruminant species did not demonstrate such a relationship on an interspecific level (Bro-Jørgensen, 2008), even though intraspecific analyses confirmed the pattern (Ashton, Tracy & de Queiroz, 2000; Meiri & Dayan, 2003). Our analyses revealed that Bergmann’s rule also applies to ruminants, if one focuses

on non-African species only (i.e. mostly non-tropical species), and only if the phylogenetic relationships between the species are considered. This finding represents one of the rarer cases where a nonsignificant finding in ordinary regression analysis becomes significant when phylogenetic relationships are accounted for (cf. fig. 1, p. 147 in Baker, 2002; Müller *et al.*, 2011a). Although testing of Bergmann’s rule was not a primary aim of the present study, this analysis represents an exquisite example of why controlling for phylogeny in comparative analyses is highly relevant.

Another ecological rule, Rensch’s rule (Abouheif & Fairbairn, 1997), indicating that the degree of sexual dimorphism among species increases with their mean body mass, was also confirmed in the present study. This rule had been confirmed previously for wild (Weckerly, 1998; but see Bro-Jørgensen, 2008) and domestic ruminants (Polák & Frynta, 2009, 2010), but not among the larger group of ungulates (Alexander *et al.*, 1979; Abouheif & Fairbairn, 1997), emphasizing the importance of testing patterns at various levels of taxonomy (Clauss, Kaiser & Hummel, 2008). The relationship between latitude and sexual dimorphism, such as postulated by Geist (hypothesis 10 in Geist, 1974b) or Isaac (2005), only tended towards significance in the overall dataset when accounting for phylogeny, and was not evident in the African or non-African subsamples (Table 1). Geist (1974b) himself provided several exceptions to his hypothesis. Elucidating reasons for sexual dimorphism in ruminants will require more variables than provided in our study.

Conflicting results have been generated on a general increase in the proportion of grass in the natural diet with body mass in various studies, and results may well depend on the selection of species at hand (reviewed in Clauss *et al.*, 2008). The finding that the relationship was not significant considering phylogenetic relationships among African species supports the concept that the African biome requires, because its C4 grasses are different from other forages on a global level, particular adaptations in grazing species (Codron *et al.*, 2008b; Codron & Clauss, 2010); these adaptations are not restricted to body mass alone and may be reflected in phylogenetic lineages. Sexual size dimorphism has been related to the type of habitat (closed/open), with dimorphic species usually occurring in more open habitats (Bro-Jørgensen, 2008). Open habitats are also generally associated with

Table 8. Linear regression analysis (in ordinary least squares, OLS, and phylogenetic generalized least squares, PGLS; differences to OLS indicated by grey shading) according to $y = a + bx$ with the day length at conception in captivity as the dependent variable, and the day length at conception in the wild as the independent variable (for species selection, see online Appendix S2)

	OLS			PGLS		
	<i>a</i>	<i>b</i>	<i>R</i> ²	<i>a</i>	<i>b</i>	<i>R</i> ²
All species (<i>N</i> = 28)	5.37 (2.56; 8.18)	0.62 (0.40; 0.83)	0.56	0.43 (−3.74; 4.60)	0.93 (0.62; 1.24)	0.56
Without five outliers (<i>N</i> = 23)	0.30 (−1.98; 2.58)	1.01 (0.83; 1.19)	0.87	1.37 (−0.55; 3.29)	0.86 (0.72; 1.00)	0.87

Results are given with 95% confidence intervals in parentheses. See Fig. 9 for the pattern of the original data and the identification of outliers.




Fig. 10. Comparison of the day length (including civil twilight) at different days of the year at the average latitude of the zoological institutions of this study (captive, 45° north), and the latitude of the free-ranging populations of (A) moose (*Alces alces*), (B) roe deer (*Capreolus capreolus*), (C) Mesopotamic fallow deer (*Dama mesopotamica*), and (D) mule deer (*Odocoileus hemionus*). The beginning of reproductive activity in the wild (data from online Appendix S2) is indicated by the black dot; the time period in which 80% of all offspring are conceived in captivity is indicated by the grey circles. Note that the different cases of time shifts between the wild and captivity as exemplified in Fig. 1 actually occur.




Fig. 11. Proportion of births (in % of all births) in two populations of goitered gazelles (*Gazella subgutturosa*) kept at a latitude of approximately 23° (Arabia) and 45° north (data used herein). Note the earlier birth peak in the population kept at lower latitude.

grazing (Pérez-Barbería, Gordon & Nores, 2001). The relationship between % grass and body mass and dimorphism might therefore reflect a habitat influence (Pérez-Barbería, Gordon & Pagel, 2002) rather than an influence of the feeding niche itself—in other words, % grass might just represent a proxy for habitat when tested against body mass. The fact

that there is no relationship between % grass and sexual dimorphism among African species suggests the interpretation that in the African biome, the browser/grazer spectrum might not be as distinctively distributed between habitats, with several savannah-based browsers and mixed feeders, and/or that in the African biome, an adaptation to grazing itself (rather than to habitat characteristics for which grazing is a proxy) may be more relevant due to the special nature of C4 grasses as mentioned above.

(2) Quantifying seasonality

In this study, we expressed the degree of seasonality as the number of succeeding 5-day intervals within a year where a certain percentage (i.e. 50, 70, 80 or 90%) of all births took place. Many previous studies did not apply a quantitative measure to the degree of reproductive seasonality, despite using quantitative data and statistical tests to allocate species to seasonality categories (Pelt, 1967). Variations in the percentages of required births result in different extents of these birth-windows (from 2 months for BPB50 up to 11 months for BPB90). The correlation between these BPB and the more traditional seasonality categories did not indicate that a certain BPB was more suited for

Table 9. Results from general linear models (retaining only variables/factors significant using original least squares, OLS) testing for the effect of reproductive seasonality (measured as the birth peak breadth in which 80% of all births occurred, BPB80), the mother-young relationship (hider/follower) or the sexual size dimorphism on the relationship between body mass and the length of the gestation period in the complete dataset ($N = 110$ species), the African ($N = 47$) and the non-African species ($N = 63$) using OLS and phylogenetic generalized least squares (PGLS; differences to OLS indicated by grey shading)

	Model	R^2	Body mass	BPB80	Hider/follower	Dimorphism
All species	OLS	0.68	$F = 200.723$ $P < 0.001$	$F = 5.035$ $P = 0.027$	$F = 10.322$ $P = 0.002$	$F = 22.877$ $P < 0.001$
	PGLS	0.50	$t = 11.000$ $P < 0.001$	$t = 3.667$ $P < 0.001$	$t = 1.000$ $P = 0.320$	$t = 2.000$ $P = 0.048$
African species	OLS	0.72	$F = 105.064$ $P < 0.001$	$F = 6.265$ $P = 0.016$	—	$F = 10.637$ $P = 0.002$
	PGLS	0.55	$t = 6.000$ $P < 0.001$	$t = 2.556$ $P = 0.014$	—	$t = 1.000$ $P = 0.323$
Non-African species	OLS	0.66	$F = 100.298$ $P < 0.001$	$F = 4.086$ $P = 0.048$	$F = 14.222$ $P < 0.001$	$F = 9.276$ $P = 0.003$
	PGLS	0.51	$t = 7.437$ $P < 0.001$	$t = 1.524$ $P = 0.133$	$t = 2.097$ $P = 0.040$	$t = 2.111$ $P = 0.039$
Non-African species (without roe deer)	OLS	0.73	$F = 129.629$ $P < 0.001$	$F = 9.018$ $P = 0.004$	$F = 15.030$ $P < 0.001$	$F = 8.843$ $P = 0.004$
	PGLS	0.59	$t = 8.399$ $P < 0.001$	$t = 2.188$ $P = 0.033$	$t = 0.954$ $P = 0.344$	$t = 1.703$ $P = 0.094$

the description of seasonality than another. Caughley & Caughley (1974) used a nonlinear regression equation to describe the birth pattern of free-ranging impala (*Aepyceros melampus*) and estimated the peak of the birth season as the median date of births (explicitly assuming a normal distribution of births around this time point). Because of the non-symmetric appearance of many birth distribution patterns (see Fig. 2), the standard deviation used in their study to describe the duration of the birthing season does not necessarily describe the real birth season very precisely. Similarly, the calculation of the quantitative reproductive asynchrony index (RAI) introduced by Kiltie (1988), or the approach of Ogutu *et al.* (2010) to describe the birth peak and the degree of birth synchrony by fitting Poisson regression models, are based on the assumption of a normal distribution around a birth peak. By contrast, the BPB, with its approach to find the smallest number of 5-day intervals in which a certain percentage of all births occurred (possible because of the high resolution of observations in captivity that are made on a daily, not a monthly basis) achieves a higher level of precision in describing the peak birth season. Rutberg (1987) and Meng *et al.* (2003) defined the birth season of a species as number of consecutive days during which 75–80% of the year's offspring are born, beginning from the first birth event that was observed in that year until 75 or 80% were achieved. Note the better representation of the real reproductive pattern of the BPB-approach (Fig. 3), because the BPB is not linked to the first birth event of the year but to the time period with the most concentrated number of birth events.

(3) Evolving seasonality

Rutberg (1987) found, using a nested analysis of variance, a high degree of variation in the birth season length in

ruminant taxa, suggesting that this characteristic is subject to a high degree of adaptive selection. Analyses on a higher taxonomic level suggest that it was the expansion into higher latitudes, and the evolution of a more seasonal reproductive pattern, that led to an increase in the overall range covered by a taxonomic ruminant subgroup (Fig. 6). In agreement with these findings, Jabbour *et al.* (1997) suggest that deer evolved in subtropical climates, and that expansion into higher latitudes was accompanied by the evolution of seasonal reproduction, and by changes in body mass (Geist, 1987). Non-seasonal reproduction is thus considered the ancestral state, as supported by our analyses for all ruminants in general; note that this does not mean that seasonality cannot evolve in low latitudes. One factor supporting ruminant radiation into higher latitudes could be that, aside from the giraffids, their gestation periods are below 1 year and can be comparatively short (*cf.* the gestation period of domestic cattle at about 278 days compared with that of similar-sized domestic horses at about 340 days - Jafar, Chapman & Casida, 1950; Bos & Van der Mey, 1980), which makes an adaptation to seasonal environments feasible without loss of a vegetation period for the interbirth interval (Kiltie, 1988). As stated in Section I, the fact that species with gestation periods longer than a seasonal cycle live or lived at higher latitudes indicates that while a short gestation period may be an advantage at higher latitudes, it is not a precondition for expansion into such regions. It might be particularly interesting to investigate the climatic conditions, the geographic distribution and the competitive replacement of fossil giraffids (Solounias, 2007) in this respect—the only true ruminants with exceptionally long gestation periods.

The finding that New World deer (Capreolini) show a numerically higher reproductive seasonality than Old world deer (Cervini) corresponds to Jabbour *et al.*'s (1997) claim that




Fig. 12. Relationship between (A) body mass and the length of the gestation period for ruminant species of different seasonality classes [in terms of the length of the birth peak breadth in which 80% of all births occur (BPB80, in days); note that for statistical evaluation, BPB80 was used as a continuous measure—see Section III.6]; (B,C) the relationship of BPB80 and the relative length of the gestation period (per unit body mass^{0.12}) in ruminant groups (B) and bovid groups (C).

due to the historical high-latitude dispersal routes of New World deer, a seasonal reproduction might be considered the ancestral state, and that tropical New World deer should therefore express seasonal reproduction when kept at higher latitudes—something not necessarily to be expected for Old World deer. Notably, several Old World deer species were amongst those in which a less seasonal pattern was

observed in captivity; however, no New World deer was amongst those considered ‘changers’, suggesting a more fixed reproductive seasonality. Nevertheless, the pampas deer (*Ozotoceros bezoarticus*), considered aseasonal in the wild (Jabbour *et al.*, 1997; category 4), had a more seasonal pattern in captivity (category 3; BPB80 < 33 days), which still corresponds to Jabbour *et al.*’s (1997) prediction.

The direction of day-length change separates the short-day (days shorter than the days before) and long-day (days longer than the days before) breeders. Ruminants are usually considered short-day breeders if seasonal, whereas many small mammals such as rodents or carnivores that conceive and give birth in the same year, and domestic horses with their longer gestation periods, are examples of long-day breeders (Sadleir, 1969; Bronson, 1989). There was only one seasonal long-day breeder in the current dataset – the Eld’s deer (*Cervus eldi*), which is in accordance with published findings on ovarian function in this species (Monfort *et al.*, 1990). For the rusa deer (*Cervus timorensis*), long-day breeding has also been reported (Van Mourik & Stelmasiak, 1990), which is in accordance with our findings—but this species was not considered seasonal in our dataset due to its long BPB80 of 225 days. All the other seasonal species were short-day breeders. The exception of the Eld’s (and rusa) deer suggests that short-day breeding need not necessarily be considered a plesiomorphic character in ruminants, but that different adjustments of breeding seasons can occur comparatively easily among closely related species. This is also demonstrated by the interesting difference between Himalayan tahr (*Hemitragus jemlahicus*) that are strictly seasonal both in captivity and the wild, and the Nilgiri tahr (*Hemitragus hyloricus*—‘the world’s most equatorial wild caprid’) that are strictly seasonal in the wild but breed throughout the year in captivity (Rice, 1988; Paré *et al.*, 1996)—an impressive example for the potential for divergence in the adaptation to seasonally fluctuating environments in closely related species.

(4) Causative factors for seasonality: interspecific comparisons

(a) Latitude

This study confirms that on a global level, latitude is the most important factor influencing reproductive seasonality. The fact that the effect of the latitude of origin was hardly attenuated by management in captivity (which ensures a constant provision of dietary resources and a predator-free environment) gives evidence for a genetically fixed reproductive pattern that is triggered by factors other than nutritional status—most likely, by photoperiod (Bradshaw & Holzapfel, 2007). The reproductive periods (conception and birth) become shorter, i.e. ruminant species become more seasonal, with increasing latitude of geographic origin of the species (and not the animals’ actual geographic position at the zoo). Our findings confirm previous qualitative and quantitative evaluations in this respect (Rutberg, 1987; Bronson, 1989; Santiago-Moreno *et al.*, 2006), with a much larger number of species.

With increasing latitude, spring starts later, but winter starts earlier in the year (Sparks & Menzel, 2002), shortening the time of favourable environmental conditions for the offspring. In Northern mammals, survival rates of newborns that are born late in the season are reduced, as they fail to accrue body reserves necessary to survive harsh winters (Clutton-Brock, Guinness & Albon, 1983; Rutberg, 1987; Festa-Bianchet, 1988; O'Donoghue & Boutin, 1995; Santiago-Moreno *et al.*, 2006). Thus, there is great selective pressure on early births and short birthing periods.

Fixed seasonal patterns of reproduction do not seem to evolve to the same extent in latitudes that have less drastic changes in resource availability and a higher degree of climatic unpredictability (tropics and sub-tropics). A possible explanation for a low proportion of seasonally reproducing species around the equator is that a limited reproductive period bears a potential threat for the development of a population. Unpredictable events, like droughts or unusual cold and wet meteorological conditions might eradicate the majority of the offspring of a whole reproductive cycle.

(b) Altitude

Although not tested in this study due to a lack of sufficient information, an additional factor that may influence the degree of reproductive seasonality is the altitude of a species' origin. Conditions at high altitudes are in many ways analogous to those at high latitudes, and it has been suggested that high-altitude habitats, such as Tibet, could have pre-adapted species to life at lower altitudes but higher latitudes (Deng *et al.*, 2011). As seasons of favourable conditions become shorter with increasing altitude, one can predict that species living at higher altitudes should show a higher degree of seasonality (Mooney & Billings, 1961; Sadleir, 1969; Körner, 2007), in addition to the seasonality due to their latitude of origin. Data on lambing periods from 30 populations of North American mountain sheep showed that the correlations between the start of the reproductive season and their duration with latitude are enhanced when latitude is replaced by a phenological index that additionally incorporates information on altitude (Bunnell, 1982). Differences in the birthing period between white-tailed deer (*Odocoileus virginianus*) and mule deer (*Odocoileus hemionus*) populations at similar latitudes, but at different altitudes, also suggests such an effect (Ransom, 1966; Bowyer, 1991). The generally short BPs of species of the Caprinae subfamily—goats and sheep, which are usually alpine species—support this hypothesis (Table 2).

(c) Predator avoidance

In studies of seasonal reproduction of ruminants, the two major explanatory scenarios usually evoked are that of an adaptation to seasonally variable resources and that of predator avoidance (Rutberg, 1987 and many publications referring to him). As outlined in Section I, follower species should benefit from a high degree of birth synchrony, whereas

hider species should benefit from asynchronous births (Ims, 1990a, b) to avoid predation.

Testing this concept by using the hider-follower dichotomy may be problematic in itself, because this dichotomy may not adequately reflect the real and potentially relevant range of behavioural adaptations across species, where for example the duration of the 'hider' stage, and the distance between mother and hidden neonate, may vary systematically (Ralls, Kranz & Lundrigan, 1986)—but a finer-scale classification of ruminant species across the hider-follower spectrum is lacking so far. Parallels in the characteristics of ungulates and macropods that correlate with the respective hider-follower strategies (Fisher, Blomberg & Owens, 2002) suggest that seasonality is not the major factor influencing the evolution of the mother-neonate relationship. Most importantly, however, the requirements of adaptation to seasonal resource availability might overrule the adaptive value of adjusting reproductive patterns to predation pressure.

Correspondingly, we did not detect an influence of the hider-follower dichotomy in the overall dataset, or in the non-African species (Table 5). Predation avoidance has been rejected as a cause for birth synchronicity in several wild ruminant species from higher latitudes (Rutberg, 1984; Festa-Bianchet, 1988; Rachlow & Bowyer, 1991; Green & Rothstein, 1993; Aanes & Andersen, 1996; Bowyer, Van Ballenberghe & Kie, 1998; Linnell & Andersen, 1998; Post *et al.*, 2003). Several of the temperate-zone ruminants are hider species yet show a distinct seasonal reproduction. Based on theoretical modelling, Ims (1990a) demonstrated that predator saturation due to birth synchrony could only work against highly specialized predators that do not switch prey species; against generalist predators (such as those preying on most hider species), the strategy is, by contrast, disadvantageous. Panzacchi *et al.* (2008) actually demonstrated such an increased predation of foxes (*Vulpes vulpes*) on roe deer fawns and concluded that birth synchrony could not have evolved primarily as an anti-predator strategy in this hider species. By contrast, Testa (2002) suggested that birth synchrony could also evolve in species susceptible to this kind of predation, because individuals born at the very beginning of a birthing period should have a higher chance of survival, due to a time lag between the very beginning of the birthing period and the reaction of predators to search for neonate prey. We conclude that even if effects of birth synchrony and timing of birth on predation risk can be demonstrated in temperate-zone ruminants (see Section I), these effects can only be considered of secondary importance at best for reproductive seasonality.

Notably, support for the anti-predation hypothesis derives from the African biome (Estes, 1976; Sinclair *et al.*, 2000b; see also Moe *et al.*, 2007, who consider the absence of reproductive seasonality as an anti-predator strategy in hider species). Among the African species, seasonal variability in available resources that can be directly linked to photoperiodism is less distinct, and therefore, the effect of predator avoidance on reproductive patterns might be more prominent. Our analysis supported the hypothesis that

those ruminants classified as followers show a higher degree of reproductive seasonality than those classified as hider species; including the effect of the hider-follower classification even led to nonsignificance in the latitude-BPB-relationship in PGLS (Table 5). Therefore, we must conclude that in the African ruminant guild, some species classified as followers show adaptations, putatively for predator swamping, that are genetically fixed (and most likely linked to photoperiodism) so that they are apparent even under captive conditions of permanent food supply and protection from predators. It is in particular in comparison to seasonal reproduction triggered by resource availability (see Section IV.5b), which is also prominent in tropical and sub-tropical regions, that the high relevance of predator avoidance for these few species must be emphasized. Yet we cannot discard the hypothesis that those African species classified as followers (*S. caffer*—the only species in this group classified as rather non-seasonal, *D. lunatus*, *D. pygargus*, *C. taurinus*, *C. gnou*, *C. nubiana*, *A. lervia*) share some other characteristics that favoured the evolution of photoperiod-linked reproductive seasonality, which would make the effect of the hider-follower dichotomy a spurious finding. Adaptation to altitude (in the Caprinae) and to photoperiod-linked seasonal rainfall and migration are possible alternative explanations in this respect. We conclude that, globally, birth synchrony should rather be understood as an adaptation to the seasonal availability of resources and not primarily as a predator-avoidance strategy in ruminants.

(d) Natural diet

Putative differences in the degree of seasonality between grazers and browsers due to a shorter flush period of grasses and the associated shorter time span of high-quality food to raise offspring when compared to browse are already controversial in the literature (Leuthold & Leuthold, 1975; Rutberg, 1987; Kiltie, 1988; Skinner *et al.*, 2002). There is no systematic effect of latitude on the occurrence of feeding types in ruminants (Table 1). In the temperate zone, seasonal variation in resource availability affects grazers and browsers alike. In the African biome, there was also no effect of feeding type, which we propose is due to the fact that at times of vegetation growth, not only grass but also browse forage is of higher nutritional value (Owen-Smith & Cooper, 1989; Meissner, Zacharias & O'Reagain, 1999).

(e) Mating type

Isaac (2005) hypothesized that latitude influenced the occurrence of sexual size dimorphism in mammals due to variations in seasonal food availability, which, under pronounced seasonal conditions, might favour the seasonal aggregation of animals and thus polygamous mating strategies. She suggested that studies that investigate sexual size dimorphism in mammals without accounting for the effect of latitude, seasonal variation or adaptations to seasonality might be 'fundamentally flawed'. There was a trend for a correlation between latitude and sexual size dimorphism in our dataset in PGLS (Table 1), and sexual size dimorphism and mating

type did show some relationship with the degree of reproductive seasonality. On the one hand, the mating type of species had an influence on the degree of reproductive seasonality as assessed by BPB50 (i.e. only considering the highest birth peak), with polygamous species having smaller birth peaks for their latitude than monogamous species. On the other hand, sexual size dimorphism was always either marginally significant or close to significant as a covariable in the general linear models applied (Table 7). We suggest that mating type and dimorphism are characteristics that follow the degree of seasonality of ruminant species, rather than facilitate it. Monogamous males that live in close proximity to their (single) partner throughout the year do not have to cope with the problem of finding the female when it is in oestrus. By contrast, males of polygamous species attempt to copulate with as many females as possible, resulting in higher sexual competition amongst the males. Such a mating strategy can more easily evolve under conditions when most females are in oestrus at the same time (such as under seasonal conditions), limiting the time period of high reproductive investment for the males. Nevertheless, many polygamous species (e.g. members of the genera *Kobus*, *Hippotragus*, and *Gazella*) demonstrate an aseasonal mode of reproduction.

(5) Environmental correlates of seasonality: intraspecific comparisons

In general, the comparison of reproductive seasonality between captive and wild populations emphasizes the high relevance of the latitude of origin, and hence photoperiodism, and the comparatively low relevance of resource-induced seasonality in the whole ruminant dataset. In contrast to the expectation that a comparison of free-ranging and captive populations might reveal a high number of species that switch to a less distinct seasonality under constant resource provision in captivity (Clauss, Hume & Hummel, 2010), the number of such species was actually remarkably low.

(a) Photoperiodism

As there was only one species that changed from a clear seasonal pattern in the wild to a clear non-seasonal pattern (category 5) in captivity (*Hemitragus hyloricus*), it can be argued that such a complete reversal of reproductive seasonality is rarely possible. Consistent with this, the ancestors of the most important domestic ruminant—cattle (*Bos taurus*, *B. indicus*) originate from populations of wild aurochs (*B. primigenius*) (Beja-Pereira *et al.*, 2006), for which - according to the latitude where their domestication started (India) - a rather aseasonal reproductive behaviour can be proposed. In fact, a similar argument can be made for some non-ruminants as well, such as the domestic pig (*Sus scrofa domestica*), which originated from populations of wild boar (*Sus scrofa*) initially in the Middle East (Giuffra *et al.*, 2000). By contrast, the ancestors of the more seasonal domestic goats (*Capra hircus*) and sheep (*Ovis aries*) had been domesticated in the highlands of the Middle East region with a more seasonal environment (Zeder & Hesse, 2000; Pedrosa *et al.*, 2005). Similarly, Ortavant

et al. (1985) stated that the original site of domestication determines the pattern of seasonal reproduction in domestic sheep as well as horse breeds.

The finding that the timing of seasonal reproduction is shifted by half a year if a species is kept at the hemisphere from which it does not originate, like *Damaliscus lunatus* and *Connochaetes gnou* in this study (and many similar observations referenced in Section I) is evidence for a general effect of day-length change on reproductive seasonality in ruminants. Further support for an effect of day-length change comes from the shift in the reproductive period with latitude (Figs 10 and 11) [see also results on *Hemitragus jemlahicus* by Paré *et al.* (1996), disregarding one outlier population out of seven], which also corresponds to other observations on individual species referenced in Section I. Finally, the finding of a high correlation between day length at conception in the wild and in captivity (Fig. 9) is strong evidence for a genetically fixed photosensitivity in ruminants that acts not only at the level of day-length changes (short-day or long-day breeders and refractoriness), but also at the level of absolute day length. To our knowledge, such strict dependence on absolute day length has not been demonstrated previously in ruminants, and is mostly considered a feature of arthropods and comparatively short-lived (and hence small) vertebrates (Bradshaw & Holzapfel, 2007). Experimental manipulation of reproductive activity in ruminants by artificial control of day length was usually not performed with the aim of detecting a specific absolute day length that triggers reproduction, but to detect patterns according to day-length change and refractoriness to long or short days (see Verme & Ozoga, 1987, for a typical example). Therefore, the result that reproductive activity is probably triggered by absolute day length in many seasonal wild ruminant species clearly warrants corroboration by experimental studies.

(b) Resource availability

There is overwhelming evidence that body condition (especially body fat stores), or in other words resource availability and the opportunity to accumulate fat reserves over a certain time period, will determine reproductive potential (Sadleir, 1969; Bronson, 1989). High resource availability and good body condition leads to earlier conceptions and births even in seasonally reproducing species (examples in Ogutu *et al.*, 2010; also see Section I). Our analysis of day length at the time of conception in free-ranging and captive populations (Fig. 9) yielded, as predicted, a numerically positive intercept, suggestive of slightly earlier conceptions in captivity - as would be expected because of the consistent provision with food. However, this effect was not significantly greater than zero, emphasizing that the influence of nutrition on seasonal breeders, though present in individual studies, is not strong enough to cause a systematic shift towards earlier conceptions and births in captivity. Again, more controlled experiments with defined day-length changes and defined levels of food provision would be required to accurately separate these effects.

When searching for triggers of reproductive seasonality, it is important to separate ultimate causes, such as favourable conditions at the time of parturition, from the proximate causes, which must be sought around the time of conception. Even though gestation period length can be adjusted to a certain degree according to actual climatic or resource conditions, the timing of conception—and hence, the timing of oestrus and rut—are the major proximate determinants of the timing of the birthing season (Ogutu *et al.*, 2010; Clements *et al.*, 2011). When investigating the effect of resource availability on birthing patterns, it is therefore important to consider these conditions at the time of conception—in ruminants, this means that the resource conditions of the previous season have to be linked to birthing patterns.

Several authors could demonstrate a correlation between rainfall, animal condition or other indicators of resource availability during the time of conception with the timing of either rutting events or birth patterns (Estes, 1976; Adams & Dale, 1998; Post, 2003; Moe *et al.*, 2007; Ryan, Knechtel & Getz, 2007; Ogutu *et al.*, 2010, 2011; Burthe *et al.*, 2011). Sometimes, confusion can occur as to whether authors refer to the birthing pattern coinciding with the rainfall pattern of the same year of the births, or of the preceding year. One consequence of resource-induced seasonality is that birth synchrony is higher after seasons with high rainfall, because a higher number of females will come into oestrus at the same time (Ogutu *et al.*, 2010). Another consequence is that reproductive seasonality may vary within a species, depending on the degree of resource seasonality different populations of that species are exposed to—either as compared between different free-ranging populations (Moe *et al.*, 2007) or as compared between free-ranging and captive populations (Pelt, 1967; Skinner *et al.*, 2002; Piening Schuler *et al.*, 2009; present study).

Periodic reproduction triggered by resources, and in particular rainfall events, is considered the least understood phenomenon in mammalian seasonality (Bronson, 2009). This may partly be due to the confusing finding that species differ in the timing of birthing periods in relation to the more or less regular rainfall events of the same season. By contrast, we suggest that the lack of a relevant pattern would probably disappear if one considered not the putative ultimate reason—births close to rainfall for maximum resource availability—but the probable proximate reason—conception at the time of optimal body condition, which should be close to the preceding season's rainfall, together with the gestation period of the species.

Several examples can be used to support this view. Hall-Martin, Skinner & Van Dyk (1975) found a conception peak in giraffe at the time of optimal nutritional status of females, whereas the resulting timing of birth did not provide newborns with optimal resources. Ryan *et al.* (2007) suggested that the comparatively long gestation period of African buffalo (*Syncerus caffer*) might be an adaptation to seasonality in the species' environment, leading, after an induction of oestrus by rainfall, to a birth of neonates close to the beginning of the next season's rainfall. On the one

hand, our own analysis suggests that the gestation period of African buffalo need not be considered peculiar due to the species low degree of reproductive seasonality; on the other hand, the authors themselves note that due to this long gestation period, African buffalo need to conceive very soon after parturition if they do not want to lose out on the next favourable resource cycle. Other species, in which oestrus, rut and conception are triggered by rainfall, might have shorter gestation periods (due to smaller body size), might give birth to their young somewhat earlier in relation to the next rainfall period, and might have more time between parturition and the time when the next conception needs to occur in order to make use of the next resource cycle. The fact that good body condition due to rainfall will trigger conception in many sympatric species at the same time will lead to a variation in birth peaks in relation to the following season's rainfall - because of the interspecific variation in gestation length. This could be a parsimonious explanation for the observation that sympatric cervid species in Nepal (Asher *et al.*, 1999) or sympatric ungulates in Southern Zimbabwe (Dasmann & Mossman, 1962) had very distinct birthing seasons (although their neonates should all thrive best at a similar time of year). It could also explain why smaller grazing species tended to give birth earlier (close to the peak protein concentration of young grasses) and larger species later (closer to the peak grass biomass) in the study of Sinclair *et al.* (2000b). These authors also wondered why oribi (*Ourebia ourebi*), although being a very small grazer, had a birth peak not around the peak of protein concentration in young grass (as the authors would expect based on their nutritional concept), but rather at the time of peak grass biomass, and speculated that this might be an adaptation to the necessity for hiding young in tall grasses. A more parsimonious explanation could be that conception might be triggered by young grass growth and that therefore the parturition peak occurs at the time of peak grass biomass - 210 days later (the oribi's gestation period; Jones *et al.*, 2009).

Recently, differences in the seasonal reproduction of sympatric Axis deer (*Axis axis*, seasonal birth peak) and gaur (*Bos gaurus*, no seasonal birth peak) in India have been explained as an adaptation to the higher nutritional requirements of the smaller species (Ahrestani *et al.*, 2011)—an ultimate reason. The proximate reason might simply be that the Axis deer have synchronized births triggered by monsoon rainfall; because of this species' shorter gestation period, enough time elapses after birth until the next period of high resources for the next conception, so that a seasonal pattern can emerge. The longer gestation periods of gaur are more likely to prevent conception at the time of high resources after a previous conception triggered by these events, which will by necessity lead to a more even spread of birth events across seasons. Reduced nutritional requirements of larger species, if existent, could act as a permissive factor that does not restrict conception to a certain time period (Ahrestani *et al.*, 2011); however, our finding that body mass did not significantly contribute to the relationship between seasonality and latitude, neither in the global

analysis nor among African or non-African species (Table 4), does not suggest that permissive factors due to systematic changes in nutritional requirements with body mass are relevant for the evolution of reproductive seasonality.

Further insight into these phenomena could be generated if not only the length of the rainfall and vegetative periods, but also rutting and the body condition of the respective species over the course of these periods was monitored, to test whether species-specific adaptations in the timing of the onset of rut exist. Additionally, it would be interesting to compare various ruminant habitats with known rainfall and resource patterns, testing whether ruminant species composition in these habitats is determined by the relationship of gestation period (as mainly determined by body mass) to the lag periods between two periods of high resource availability. It could be hypothesized that species are favoured whose gestation period allows an optimal use of the respective periods of high resource availability without losing reproductive potential due to delays between parturition and the following conception (Kiltie, 1988).

Conceptually, one might argue that the resource-dependence of conception did not so much evolve as a response to a predictive trigger—in the sense that this year's rainfall might also indicate next year's rainfall -, but rather as a mechanism of reducing the energetic demands of reproduction during times of scarce resources. Note that the latter interpretation removes the element of prediction from resource-dependent reproduction, but shifts the focus to an 'emergency break' function that prevents parents from fatally depleting their own resources. Such a mechanism has been proposed previously by Owen-Smith (1988, p. 184) for white rhinoceros (*Ceratotherium simum*); he found evidence that high planes of nutrition were not so much stimulating reproduction, but that rather low planes of nutrition prevented it; similar findings on elephants are also provided by this author. Synchronised seasonality then simply ensues because after a period of scarcity, many adults will enter reproductive activity in the next season of high resource availability simultaneously. This view is also more compatible with current understanding of the hormonal modulation of reproduction by signals referring to body condition such as leptin (e.g. Zieba *et al.*, 2007).

(c) Specific examples

The peculiar case of the springbok (*Antidorcas marsupialis*), the only species that changed to a more seasonal birth pattern in captivity in this study, has already been noted by other authors for zoos in both hemispheres (Pelt, 1967; Skinner *et al.*, 2002). To date, a similar change of birth seasonality was only observed in domestic hoofstock from the tropics—i.e. Creole goats - which have a rather unseasonal reproductive pattern in the tropics, but demonstrate a marked seasonality when subjected to the large photoperiodic variation in the northern hemisphere (Chemineau *et al.*, 2004). Skinner *et al.* (1996) observed irregular rutting in springbok males but noted that the majority of ruts occurred closely to

the beginning of the long-day period after the southern-hemisphere winter solstice in June; by contrast, conception in our zoo population was concentrated during the short-day period of the northern-hemisphere year. Reasons for this discrepancy remain unclear, but could lie in the fact that the animals observed in the wild did not respond to photoperiodism but to resource cues. Later Skinner *et al.* (2001) observed spontaneous ovarian cycling in recently matured springbok ewes from November to June, i.e. in the short-day period of the southern hemisphere; the animals ceased cycling close to the summer solstice and started cycling again, after an interruption of about 4 months, during the following short-day period. These findings are easier to reconcile with the observations from captivity, and suggest that in the wild, there is a strong modulation of seasonal reproduction by resource availability, leading to variable conception dates. In captivity, because resources are continually provided, the photoperiodic signal is less modulated and hence seasonality more readily detected.

By contrast, the striking difference in seasonality between the free-ranging and the captive Nilgiri tahr (*Hemitragus hyloricus*) populations, with non-seasonal reproduction in captivity, do not support the hypothesis that reproduction is linked to photoperiod in this species. Rice (1988) suggested changes in photoperiod due to the monsoon (with short days due to clouded skies) as the proximate causes of seasonal reproduction in free-ranging Nilgiri tahr, but this should then also have an effect in temperate-zone zoos. In this species, reproduction is therefore probably only resource-constrained.

(6) Gestation length and reproductive seasonality

Several authors have noted the relevance of gestation period lengths for seasonal reproduction (Kiltie, 1988; Owen-Smith, 1988; Jabbour *et al.*, 1997). In particular, it has been suggested that smaller species might be forced to evolve particularly long gestation periods, in order to bridge the time between favourable rutting conditions and favourable birthing conditions (Jabbour *et al.*, 1997), and that large animals need to shorten their gestation period so that both parturition and the subsequent conception can both occur in the same (short) vegetation period; otherwise, they would lose one seasonal cycle for reproduction (Kiltie, 1988). The results of our study clearly support the latter concept, in that more seasonal ruminants have comparatively shorter gestation periods (Fig. 12). In contrast, there was no evidence for a systematic increase in gestation period with decreasing body mass in seasonal ruminants (i.e. the seasonality-body mass interaction was not significant when investigating the influence of both factors on gestation length).

It needs to be acknowledged that the concept of one assumed fixed gestation period length per species evidently does not reflect reality. The length of the gestation period can change, within a species, with parity (primiparous *versus* multiparous) and with the condition of the mother (and hence with variable environmental conditions), with the sex and the number of the offspring, and with conception date itself

(reviewed in Clements *et al.*, 2011). While such fine-tuning of gestation periods does occur, it is nevertheless the timing of conception—and hence, the average gestation period of a species—that is the major determinant of the timing of the birthing season (Ogutu *et al.*, 2010; Clements *et al.*, 2011).

As seasonal environments have short periods of sufficient food supply, these time periods have to be used to full capacity by resident species. The mating season is often accompanied by a decreased food intake not only due to lack of time (Geist, 1974a; Lincoln & Short, 1980) but probably also as a side-effect of physiological processes associated with chemical cues of rut (Miquelle, 1990). It may be advantageous in seasonal environments to set this period to a time of the year when food supply is no longer at its optimum and body fat stores have accumulated maximally (Owen-Smith, 2002, pp. 156–158), so that no valuable opportunities for building up body reserves for the winter period are wasted. To achieve this goal, seasonal reproducing species of higher latitudes might shift conception and rut more to the end of the vegetative period, and hence closer to the next favourable period of parturition. It remains to be investigated whether differences between various taxonomic groups exist in the degree that gestation periods can be shortened. This characteristic may have made some ruminant groups particularly pre-adapted candidates for the occupation of seasonal habitats (in contrast to giraffids, camelids or perissodactyls) as suggested above.

On the other hand, a prolongation of gestation periods appears to be rare among ruminants. Increasing gestation length by the mechanism of delayed implantation—possibly a conserved characteristic in mammals (Ptak *et al.*, 2012)—has been documented in the roe deer (*Capreolus capreolus*) (Short & Hay, 1966). Note that the inclusion of this dramatic outlier affected the statistical results (Table 9). Delayed implantation was also suspected to occur in several other ruminant species such as the New World deer *Blastocerus* and *Hippocamelus* (not represented in this study) (Jabbour *et al.*, 1997), pudu (*Pudu puda*) (Bubenik *et al.*, 2000), and in the Père David's deer (*Elaphurus davidianus*) (Brinklow & Loudon, 1993). Our findings reinforce such suspicions insofar as both the pudu and the Père David's deer are among the more seasonal species that have, for their body size, comparatively long gestation periods; the same is true for the pronghorn antelope (*Antilocapra americana*) (Fig. 12A). In the pronghorn, foetal growth might be delayed as compared to other ungulates (fig. 1 in Robbins & Robbins, 1979), and yet unconfirmed reports of an increasing gestation period with latitude (Buechner, 1950) are also compatible with delayed embryonic development. Future studies might detect retarded embryonic growth, or delayed implantation, in these species.

V. CONCLUSIONS

(1) We expressed the degree of birth seasonality of species as a continuous variable: the time period in which a certain percentage of all offspring are born.

(2) Latitude of geographical origin of species' was found to be the only significant factor influencing the degree of birth

seasonality, i.e. the birth seasons become shorter (animal species become more seasonal) with increasing latitude of geographical origin.

(3) As only about 11% of the tested species distinctively changed their birth seasonality pattern in captivity when compared to wild populations, the birth season length in zoos can be seen as a good predictor for birth season length in the wild for temperate species.

(4) It is well recognized that the evolved adaptation of using specific photoperiodic cues for the timing of reproductive activity in temperate species can lead to problems under current global warming scenarios. In generally warmer conditions, plants will tend to emerge earlier, while reproduction cued to photosignals stays inert and thus becomes out of synch with the beginning of forage availability, with potentially deleterious effects (Post & Forchhammer, 2008; Post *et al.*, 2008; Bronson, 2009). Changes in breeding phenology, linked to variation in resource availability due to warmer climate, have been documented in both seasonal and aseasonal breeders (Burthe *et al.*, 2011; Moyes *et al.*, 2011). The degree to which ruminants will be affected by the mismatch between photoperiod and resource availability will depend on the speed at which adjustments of the photoperiodic responses can evolve—or to which degree less-seasonal species can replace more-seasonal species in their respective ecosystems.

VI. ACKNOWLEDGEMENTS

We thank the Georg and Bertha Schwyzer-Winiker-Stiftung and the Vontobel-Stiftung for financial support, the World Association of Zoos and Aquariums for enabling data transfer from ISIS, all participating zoos for their consistent data collection, Jeanne Peter for graphics support, and Norman Owen-Smith for meticulous comments on earlier manuscript versions.

VII. REFERENCES

- AANES, R. & ANDERSEN, R. (1996). The effects of sex, time of birth, and habitat on the vulnerability of roe deer fawns to red fox predation. *Canadian Journal of Zoology* **74**, 1857–1865.
- ABOUHEIF, E. & FAIRBAIRN, D. J. (1997). A comparative analysis of allometry for sexual size dimorphism: assessing Rensch's rule. *American Naturalist* **149**, 540–562.
- ADAMS, L. G. & DALE, B. W. (1998). Timing and synchrony of parturition in Alaskan caribou. *Journal of Mammalogy* **79**, 287–294.
- ADAMS, L. G., SINGER, F. J. & DALE, B. W. (1995). Caribou calf mortality in Denali National Park, Alaska. *Journal of Wildlife Management* **59**, 584–594.
- AHRESTANI, F. S., VAN LANGEVELDE, F., HEITKÖNIG, I. M. A. & PRINS, H. H. T. (2011). Contrasting timing of parturition of chital (*Axis axis*) and gaur (*Bos gaurus*) in tropical South India—the role of body mass and seasonal forage quality. *Oikos* (doi: 10.1111/j.1600-0706.2011.20244.x).
- ALEXANDER, R. D., HOOGLAND, J. L., HOWARD, R. D., NOONAN, K. M. & SHERMAN, P. W. (1979). Sexual dimorphism and breeding system in pinnipeds, ungulates, primates, and humans. In *Evolutionary Biology and Human Social Behavior: An Anthropological Perspective* (eds N. A. CHAGNON and W. IRONS), pp. 402–435. Duxbury, North Scituate.
- ASHER, G. W., MONFORT, S. L. & WEMMER, C. (1999). Comparative reproductive function in cervids: implications for management of farm and zoo populations. *Journal of Reproduction and Fertility Supplement* **54**, 143–156.
- ASHTON, K. G., TRACY, M. C. & DE QUEIROZ, A. (2000). Is Bergmann's rule valid for mammals? *American Naturalist* **156**, 390–415.
- BAKER, R. H. (2002). Comparative methods. In *Techniques in Molecular Systematics and Evolution* (eds R. DeSalle, G. Giribet and W. Wheeler), pp. 146–161. Birkhäuser Verlag, Basel.
- BARRETTE, C. (1987). The comparative behavior and ecology of Chevrotains, Musk deer, and morphologically conservative deer. In *Biology and Management of the Cervidae* (ed. C. M. Wemmer), pp. 200–213. Smithsonian Institution Press, Washington.
- BEJA-PEREIRA, A., CARAMELLI, D., LALUEZA-FOX, C., VERNESI, C., FERRAND, N., CASOLI, A., GOYACHE, F., ROYO, L. J., CONTI, S., LARI, M., MARTINI, A., OURAGH, L., MAGID, A., ATASH, A., ZSOLNAI, A., BOSCATO, P., TRIANTAPHYLIDIS, C., PLOUMI, K., SINEO, L., MALLEGNI, F., TABERLET, P., ERHARDT, G., SAMPIETRO, L., BERTRANPETIT, J., BARBUJANI, G., LUIKART, G. & BERTORELLE, G. (2006). The origin of European cattle: evidence from modern and ancient DNA. *Proceedings of the National Academy of Sciences* **103**, 8113–8118.
- BININDA-EMONDS, O. R. P., CARDILLO, M., JONES, K. E., MACPHEE, R. D. E., BECK, R. M. D., GRENYER, R., PRICE, S. A., VOS, R. A., GITTLEMAN, J. L. & PURVIS, A. (2007). The delayed rise of present-day mammals. *Nature* **446**, 507–512.
- BONENFANT, C., LOE, L. E., MYSTERUD, A., LANGVATN, R., STENSETH, N. C., GAILLARD, J.-M. & KLEIN, F. (2004). Multiple causes of sexual segregation in European red deer enlightenments from varying breeding phenology at high and low latitude. *Proceedings of the Royal Society B* **271**, 883–892.
- BOS, H. & VAN DER MEY, G. J. W. (1980). Length of gestation periods of horses and ponies belonging to different breeds. *Livestock Production Science* **7**, 181–187.
- BOWYER, R. T. (1991). Timing of parturition and lactation in southern mule deer. *Journal of Mammalogy* **72**, 138–145.
- BOWYER, R. T., VAN BALLEMBERGHE, V. & KIE, J. G. (1998). Timing and synchrony of parturition in Alaskan moose: long-term versus proximal effects of climate. *Journal of Mammalogy* **79**, 1332–1344.
- BRADSHAW, W. E. & HOLZAPFEL, C. M. (2007). Evolution of animal photoperiodism. *Annual Reviews of Ecology, Evolution, and Systematics* **38**, 1–25.
- BRAND, D. J. (1963). Records of mammals bred in the national zoological gardens of South Africa during the period 1908 to 1960. *Proceedings of the Zoological Society of London* **140**, 617–659.
- BRINKLOW, B. R. & LOUDON, A. S. (1993). Gestation period in the Père David's deer (*Elaphurus davidianus*): evidence for embryonic diapause or delayed development. *Reproduction, Fertility and Development* **5**, 567–575.
- BRO-JØRGENSEN, J. (2008). Dense habitats selecting for small body size: a comparative study on bovids. *Oikos* **117**, 729–737.
- BRONSON, F. H. (1985). Mammalian reproduction: an ecological perspective. *Biology of Reproduction* **32**, 1–26.
- BRONSON, F. H. (1989). *Mammalian Reproductive Biology*. University of Chicago Press, Chicago.
- BRONSON, F. H. (2009). Climate change and seasonal reproduction in mammals. *Philosophical Transactions of the Royal Society B* **364**, 3331–3340.
- BUBENIK, G., REYES, E., SCHAMS, D., LOBOS, A. & BARTOŠ, L. (2000). Pudu, the smallest deer of the world: 10 years of endocrine studies of Southern Pudu (*Pudu pudu*) in Chile. *Zeitschrift für Jagdwissenschaft* **46**, 129–138.
- BUECHNER, H. K. (1950). Life history, ecology, and range use of the pronghorn antelope in Trans-Pecos Texas. *American Midland Naturalist* **43**, 257–354.
- BUNNELL, F. L. (1982). The lambing period of Mountain sheep: synthesis, hypotheses, and tests. *Canadian Journal of Zoology* **60**, 1–14.
- BURTHE, S., BUTLER, A., SEARLE, K. R., HALL, S. J. G., THACKERAY, S. J. & WANLESS, S. (2011). Demographic consequences of increased winter births in a large aseasonally breeding mammal (*Bos taurus*) in response to climate change. *Journal of Animal Ecology* **80**, 1134–1144.
- CAUGHLEY, G. (1971). The season of births for northern-hemisphere ungulates in New Zealand. *Mammalia* **35**, 204–219.
- CAUGHLEY, G. & CAUGHLEY, J. (1974). Estimating median date of birth. *The Journal of Wildlife Management* **38**, 552–556.
- CHEMINEAU, P., DAVEAU, A., COGNIE, Y., AUMONT, G. & CHESNEAU, D. (2004). Seasonal ovulatory activity exists in tropical Creole female goats and Black Belly ewes subjected to a temperate photoperiod. *BMC Physiology* **4**, 12.
- CHEMINEAU, P., GUILLAUME, D., MIGAUD, M., THIÉRY, J. C., PELLICER-RUBIO, M. T. & MALPAUX, B. (2008). Seasonality of reproduction in mammals: intimate regulatory mechanisms and practical implications. *Reproduction in Domestic Animals* **43**, 40–47.
- CLAUSS, M., HUME, I. D. & HUMMEL, J. (2010). Evolutionary adaptations of ruminants and their potential relevance for modern production systems. *Animal* **4**, 979–992.
- CLAUSS, M., KAISER, T. & HUMMEL, J. (2008). The morphophysiological adaptations of browsing and grazing mammals. In *The Ecology of Browsing and Grazing* (eds I. J. Gordon and H. H. T. Prins), pp. 47–88. Springer, Heidelberg.
- CLEMENTS, M. N., CLUTTON-BROCK, T. H., ALBON, S. D., PEMBERTON, J. M. & KRUUK, L. E. B. (2011). Gestation length variation in a wild ungulate. *Functional Ecology* **25**, 691–703.
- CLUTTON-BROCK, T. H., GUINNESS, F. E. & ALBON, S. D. (1983). The costs of reproduction to Red deer hinds. *Journal of Animal Ecology* **52**, 367–383.

- CODRON, D., BRINK, J. S., ROSSOUW, L. & CLAUSS, M. (2008a). The evolution of ecological specialization in southern African ungulates: competition- or physical environmental turnover? *Oikos* **117**, 344–353.
- CODRON, D., BRINK, J. S., ROSSOUW, L., CLAUSS, M., CODRON, J., LEETHORP, J. A. & SPONHEIMER, M. (2008b). Functional differentiation of African grazing ruminants: an example of specialized adaptations to very small changes in diet. *Biological Journal of the Linnean Society* **94**, 755–764.
- CODRON, D. & CLAUSS, M. (2010). Rumen physiology constrains diet niche: linking digestive physiology and food selection across wild ruminant species. *Canadian Journal of Zoology* **88**, 1129–1138.
- CÔTÉ, S. D. & FESTA-BIANCHET, M. (2001). Birthdate, mass and survival in mountain goat kids: effects of maternal characteristics and forage quality. *Oecologia* **127**, 230–238.
- DARDENE, H. (2012). Melatonin-dependent timing of seasonal reproduction by the Pars tuberalis: pivotal roles for long daylengths and thyroid hormones. *Journal of Neuroendocrinology* **24**, 249–266.
- DASMANN, R. F. & MOSSMAN, A. S. (1962). Reproduction in some ungulates in Southern Rhodesia. *Journal of Mammalogy* **43**, 533–537.
- DAUPHINÉ, T. C. & MCCLURE, R. L. (1974). Synchronous mating in Canadian barren-ground caribou. *Journal of Wildlife Management* **38**, 54–66.
- DENG, T., WANG, X., FORTIELUS, M., LI, Q., WANG, Y., TSENG, Z. J., TAKEUCHI, G. T., SAYLOR, J. E., SÄILÄ, L. K. & XIE, G. (2011). Out of Tibet: pliocene woolly rhino suggests high-plateau origin of ice age megaherbivores. *Science* **333**, 1285.
- DITTRICH, L. (1970). Beitrag zur Fortpflanzungsbiologie afrikanischer Antilopen im Zoologischen Garten. *Der Zoologische Garten NF* **39**, 16–40.
- ESTES, R. D. (1976). The significance of breeding synchrony in the Wildebeest. *African Journal of Ecology* **14**, 135–152.
- FAIRALL, N. (1968). The reproductive season of some mammals in the Kruger National Park. *Zoologica Africana* **3**, 189–210.
- FELSENSTEIN, J. (1985). Phylogenies and the comparative method. *American Naturalist* **125**, 1–15.
- FESTA-BIANCHET, M. (1988). Birthdate and survival in Bighorn lambs (*Ovis canadensis*). *Journal of Zoology* **214**, 653–661.
- FISHER, D. O., BLOMBERG, S. P. & OWENS, I. P. F. (2002). Convergent maternal care strategies in ungulates and macropods. *Evolution* **56**, 167–176.
- FLETCHER, T. J. (1974). The timing of reproduction in Red deer (*Cervus elaphus*) in relation to latitude. *Journal of Zoology* **172**, 363–367.
- FLOOD, P. F. & TEDESCO, S. C. (1997). Relationship between conception date and latitude in muskoxen. *Rangifer* **17**, 25–30.
- FLOWER, S. S. (1932). Notes on the recent mammals of Egypt, with a list of the species recorded from that kingdom. *Proceedings of the Royal Society of London* **102**, 369–450.
- FLYDAL, K. & REIMERS, E. (2002). Relationship between calving time and physical condition in three wild Reindeer (*Rangifer tarandus*) populations in southern Norway. *Wildlife Biology* **8**, 145–151.
- FORSYTHE, W. C., RYKIEL, E. J., STAHL, R. S., WU, H.-I. & SCHOOLFIELD, R. M. (1995). A model comparison for daylength as a function of latitude and day of year. *Ecological Modelling* **80**, 87–95.
- GEIST, V. (1974a). *On the Relationship of Ecology and Behavior in the Evolution of Ungulates: Theoretical Considerations*. IUCN, Morges.
- GEIST, V. (1974b). On the relationship of social evolution and ecology in ungulates. *American Zoologist* **14**, 205–220.
- GEIST, V. (1987). On speciation in Ice Age mammals, with special reference to cervids and caprids. *Canadian Journal of Zoology* **65**, 1067–1084.
- GIUFFRÀ, E., KIJAS, J. M. H., AMARGER, V., CARLBORG, O., JEON, J. T. & ANDERSSON, L. (2000). The origin of the domestic pig: independent domestication and subsequent introgression. *Genetics* **154**, 1785–1791.
- GREEN, W. C. H. & ROTHSTEIN, A. (1993). Asynchronous parturition in bison: implications for the hider-follower-dichotomy. *Journal of Mammalogy* **74**, 920–925.
- GREGG, M. A., BRAY, M., KILBRIDE, K. M. & DUNBAR, M. R. (2001). Birth synchrony and survival of pronghorn fawns. *Journal of Wildlife Management* **65**, 19–24.
- HALL-MARTIN, A. J., SKINNER, J. D. & VAN DYK, J. M. (1975). Reproduction in the giraffe in relation to some environmental factors. *African Journal of Ecology* **13**, 237–248.
- HAMILTON, W. J. & BLAXTER, K. L. (1980). Reproduction in farmed red deer. 1. Hind and stag fertility. *Journal of Agricultural Science* **95**, 261–273.
- HAZLERIGG, D. G. & LINCOLN, G. A. (2011). Hypothesis: cyclical histogenesis is the basis of circannual timing. *Journal of Biological Rhythms* **26**, 471–485.
- HAZLERIGG, D. G. & LOUDON, A. (2008). New insights into ancient seasonal life timers. *Current Biology* **18**, R795–R804.
- HODGES, J. K., BEVAN, D. J., CELMA, M., HEARN, J. P., JONES, D. M., KLEIMAN, D. G., KNIGHT, J. A. & MOORE, H. D. M. (1984). Aspects of the reproductive endocrinology of the female Giant panda (*Ailuropoda melanoleuca*) in captivity with special reference to the detection of ovulation and pregnancy. *Journal of Zoology* **203**, 253–267.
- HUT, R. A. & BEERSMA, D. G. M. (2011). Evolution of time-keeping mechanisms: early emergence and adaptation to photoperiod. *Philosophical Transactions of the Royal Society B* **366**, 2141–2154.
- IMS, R. A. (1990a). On the adaptive value of reproductive synchrony as a predator swamping strategy. *American Naturalist* **136**, 485–498.
- IMS, R. A. (1990b). The ecology and evolution of reproductive synchrony. *Trends in Ecology and Evolution* **5**, 135–140.
- ISAAC, J. L. (2005). Potential causes and life-history consequences of sexual size dimorphism in mammals. *Mammal Review* **35**, 101–115.
- JABBOUR, H. N., HAYSEN, V. & BRUFORD, M. W. (1997). Conservation of deer: contributions from molecular biology, evolutionary ecology, and reproductive physiology. *Journal of Zoology* **243**, 461–484.
- JAFAR, S. M., CHAPMAN, A. B. & CASIDA, L. E. (1950). Causes of variation in length of gestation in dairy cattle. *Journal of Animal Science* **9**, 593–601.
- JARMAN, P. J. (1983). Mating system and sexual dimorphism in large, terrestrial, mammalian herbivores. *Biological Reviews* **58**, 485–520.
- JONES, K. E., BIELBY, J., CARDILLO, M., FRITZ, S. A., O'DELL, J., ORME, C. D. L., SAFI, K., SECHREST, W., BOAKES, E. H., CARBONE, C., CONNOLLY, C., CUTTS, M. J., FOSTER, J. K., GRENYER, R., HABIB, M., PLASTER, C. A., PRICE, S. A., RIGBY, E. A., RIST, J., TEACHER, A., BININDA-EMONDS, O. R. P., GITTLEMAN, J. L., MACE, G. M., PURVIS, A. & MICHENER, W. K. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, 2648.
- KILTIE, R. A. (1984). Seasonality, gestation time, and large mammal extinctions. In *Quaternary Extinctions: A Prehistoric Revolution* (eds P. S. MARTIN and R. G. KLEIN), pp. 299–314. University of Arizona Press, Tucson.
- KILTIE, R. A. (1988). Gestation as a constraint on the evolution of seasonal breeding in mammals. In *Evolution of Life Histories of Mammals. Theory and Pattern* (ed. M. S. BOYCE), pp. 257–289. Yale University Press, New Haven.
- KIRKWOOD, J. K., GASKIN, C. D. & MARKHAM, J. (1987). Perinatal mortality and season of birth in captive wild ungulates. *Veterinary Record* **120**, 386–390.
- KÖRNER, C. (2007). The use of 'altitude' in ecological research. *Trends in Ecology & Evolution* **22**, 569–574.
- LANE, J. E., BOUTIN, S., SPEAKMAN, J. R. & HUMPHRIES, M. M. (2010). Energetic costs of male reproduction in a scramble competition mating system. *Journal of Animal Ecology* **79**, 27–34.
- LENT, P. C. (1974). Mother-infant relationships in ungulates. In *The behavior of Ungulates and its Relation to Management* (eds V. GEIST and F. WALTHER), pp. 14–55. International Union for the Conservation of Nature and Natural Resources, Morges.
- LEUTHOLD, W. & LEUTHOLD, B. M. (1975). Temporal patterns of reproduction in ungulates of Tsavo East National Park, Kenya. *African Journal of Ecology* **13**, 159–169.
- LINCOLN, G. A. (1985). Seasonal breeding in deer. In *Biology of Deer Production* (Volume 22, eds P. F. FENNESSEY and K. R. DREW), pp. 165–180. Bulletin of the Royal Society of New Zealand, Wellington.
- LINCOLN, G. A., CLARKE, I. J., HUT, R. A. & HAZLERIGG, D. G. (2006). Characterizing a mammalian circannual pacemaker. *Science* **314**, 1941–1944.
- LINCOLN, G. A. & SHORT, R. V. (1980). Seasonal breeding: nature's contraceptive. *Recent Progress in Hormone Research* **36**, 1–53.
- LINNELL, J. D. C. & ANDERSEN, R. (1998). Timing and synchrony of birth in a hider species, the Roe deer (*Capreolus capreolus*). *Journal of Zoology* **244**, 497–504.
- LOE, L. E., BONENFANT, C., MYSTERUD, A., GAILLARD, J.-M., LANGVATN, R., KLEIN, F., CALENGE, C., ERGON, T., PETTORELLI, N. & STENSETH, N. C. (2005). Climate predictability and breeding phenology in red deer: timing and synchrony of rutting and calving in Norway and France. *Journal of Animal Ecology* **74**, 579–588.
- LOISON, A., GAILLARD, J.-M., PÉLABON, C. & YOCOZ, N. G. (1999). What factors shape sexual size dimorphism in ungulates? *Evolutionary Ecology Research* **1**, 611.
- MADDISON, W. P. & MADDISON, D. R. (2006). Mesquite: a modular system for evolutionary analysis. Available at <http://mesquiteproject.org>; accessed on 1 December 2010.
- MARSHALL, F. H. A. (1937). On the change over in the oestrous cycle in animals after transference across the equator, with further observations on the incidence of the breeding seasons and the factors controlling sexual periodicity. *Proceedings of the Royal Society of London B* **122**, 413–428.
- MARTINS, E. (2004). *COMPARE, Version 4.6. Computer Programs for the Statistical Analysis of Comparative Data*. Department of Biology, Indiana University, Bloomington, Distributed by the author at <http://compare.bio.indiana.edu/>; accessed on 1 August 2005.
- MARTINS, E. P. & HANSEN, T. F. (1997). Phylogenesis and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. *American Naturalist* **149**, 646–667.
- MASUI, M., HIRAMATSU, H., NOSE, N., NAKAZATO, R., SAGAWA, Y., TAJIMA, H. & SAITO, K. (1989). Successful artificial insemination in the Giant panda (*Ailuropoda melanoleuca*) at Ueno Zoo. *Zoo Biology* **8**, 17–26.
- MATSUBAYASHI, H., BOSI, E. & KOHSHIMA, S. (2003). Activity and habitat use of Lesser Mouse-Deer (*Tragulus javanicus*). *Journal of Mammalogy* **84**, 234–242.
- MCGINNIS, B. S. & DOWNING, R. L. (1977). Factors affecting the peak of White-tailed deer fawning in Virginia. *Journal of Wildlife Management* **41**, 715–719.
- MEIRI, S. & DAYAN, T. (2003). On the validity of Bermann's rule. *Journal of Biogeography* **30**, 331–351.

- MEISSNER, H. H., ZACHARIAS, P. J. K. & O'REAGAN, P. J. (1999). Forage quality (feed value). In *Veld Management in South Africa* (ed. N. M. TANTON), pp. 139–168. University of Natal Press, Pietermaritzburg.
- MENG, X., YANG, Q., FENG, Z., XIA, L., JIANG, Y. & WANG, P. (2003). Timing and synchrony of parturition in Alpine musk deer (*Moschus sifanicus*). *Folia Zoologica* **52**, 39–50.
- MIQUELLE, D. G. (1990). Why don't bull moose eat during the rut? *Behavioral Ecology and Sociobiology* **27**, 145–151.
- MITCHELL, B. & LINCOLN, G. A. (1973). Conception dates in relation to age and condition in two populations of red deer in Scotland. *Journal of Zoology* **171**, 141–152.
- MOE, S. R., RUTINA, L. P. & DU TOIT, J. T. (2007). Trade-off between resource seasonality and predation risk explains reproductive chronology in impala. *Journal of Zoology* **273**, 237–243.
- MONFORT, S. L., WEMMER, C. M., KEPLER, T. H., BUSH, M., BROWN, J. L. & WILDT, D. E. (1990). Monitoring ovarian function and pregnancy in Eld's deer (*Cervus eldi*) by evaluating urinary steroid metabolite excretion. *Journal of Reproduction and Fertility* **88**, 271–281.
- MONTGOMERY, G. W., SCOTT, I. C. & HUDSON, N. (1985). An interaction between season of calving and nutrition on the resumption of ovarian cycles in post-partum beef cattle. *Journal of Reproduction and Fertility* **73**, 45–50.
- MOONEY, H. A. & BILLINGS, W. D. (1961). Comparative physiological ecology of arctic and alpine populations of *Oxyria digyna*. *Ecological Monographs* **31**, 1–29.
- MOYES, K., NUSSEY, D. H., CLEMENTS, M. N., GUINNESS, F. E., MORRIS, A., MORRIS, S., PEMBERTON, J. M., KRUUK, L. E. B. & CLUTTON-BROCK, T. H. (2011). Advancing breeding phenology in response to environmental change in a wild red deer population. *Global Change Biology* **17**, 2455–2469.
- MÜLLER, D. W. H., CATON, J., CODRON, D., SCHWARM, A., LENTLE, R., STREICH, W. J., HUMMEL, J. & CLAUS, M. (2011a). Phylogenetic constraints on digesta separation: variation in fluid throughput in the digestive tract in mammalian herbivores. *Comparative Biochemistry and Physiology A* **160**, 207–220.
- MÜLLER, D. W. H., ZERBE, P., CODRON, D., CLAUS, M. & HATT, J. M. (2011b). A long life among ruminants: giraffids and other special cases. *Schweizer Archiv für Tierheilkunde* **153**, 515–519.
- NIEVERGELT, B. (1966). Unterschiede in der Setzzeit beim Alpensteinbock (*Capra ibex*). *Revue Suisse de Zoologie* **73**, 446–454.
- NIEVERGELT, B. (1974). A comparison of rutting behaviour and grouping in the Ethiopian and Alpine ibex. In *The Behavior of Ungulates and its Relation to Management* (eds V. GEIST and F. WALTHER), pp. 324–340. International Union for the Conservation of Nature and Natural Resources, Morges.
- NOWAK, R. M. (1991). Chevrotains, or mouse deer. *Walker's Mammals of the World* (Volume 2), pp. 1359–1362. Johns Hopkins University Press, Baltimore.
- O'DONOGHUE, M. & BOUTIN, S. (1995). Does reproductive synchrony affect juvenile survival rates of northern mammals? *Oikos* **74**, 115–121.
- OGUTU, J., PIEPHO, H.-P., DUBLIN, H., BHOLA, N. & REID, R. (2010). Rainfall extremes explain interannual shifts in timing and synchrony of calving in topi and warthog. *Population Ecology* **52**, 89–102.
- OGUTU, J., PIEPHO, H.-P., DUBLIN, H., BHOLA, N. & REID, R. (2011). Dynamics of births and juvenile recruitment in Mara–Serengeti ungulates in relation to climatic and land use changes. *Population Ecology* **53**, 195–213.
- ORTAVANT, R., PELLETIER, J., RAVAUULT, J. P., THIMONIER, J. & VOLLAND-NAIL, P. (1985). Photoperiod: main proximal and distal factor of the circannual cycle of reproduction in farm mammals. *Oxford Reviews of Reproductive Biology* **7**, 305–345.
- OWEN-SMITH, N. (1988). *Megaherbivores—The Influence of Very Large Body Size on Ecology*. Cambridge University Press, Cambridge.
- OWEN-SMITH, N. (2002). *Adaptive Herbivore Ecology. From Resources to Populations in Variable Environments*. Cambridge University Press, Cambridge.
- OWEN-SMITH, N. & COOPER, S. M. (1989). Nutritional ecology of a browsing ruminant, the kudu (*Tragelaphus strepsiceros*), through the seasonal cycle. *Journal of Zoology* **219**, 29–43.
- PAGEL, M. (1999). Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884.
- PANZACCHI, M., LINNELL, J. D. C., ODDEN, J., ODDEN, M. & ANDERSEN, R. (2008). When a generalist becomes a specialist: patterns of red fox predation on roe deer fawns under contrasting conditions. *Canadian Journal of Zoology* **86**, 116–126.
- PARÉ, P., BARRETTE, C. & PRESCOTT, J. (1996). Seasonal reproduction of captive Himalayan tahr (*Hemitragus jemlahicus*) in relation to latitude. *Journal of Mammalogy* **77**, 826–832.
- PEDROSA, S., UZUN, M., ARRANZ, J.-J., GUTIÉRREZ-GIL, B., SAN PRIMITIVO, F. & BAYÓN, Y. (2005). Evidence of three maternal lineages in near eastern sheep supporting multiple domestication events. *Proceedings of the Royal Society B: Biological Sciences* **272**, 2211–2217.
- PELT, F. L. (1967). Remarks on breeding seasons of some *Artiodactyla* in captivity. *Der Zoologische Garten NF* **34**, 293–296.
- PENFOLD, L. M. (2005). Reproductive physiology and artificial insemination studies in wild and captive Gerenuk (*Litocranius walleri walleri*). *Reproduction, Fertility and Development* **17**, 707.
- PÉREZ-BARBERÍA, F. J., GORDON, I. J. & NORES, C. (2001). Evolutionary transitions among feeding styles and habitats in ungulates. *Evolutionary Ecology Research* **3**, 221–230.
- PÉREZ-BARBERÍA, F. J., GORDON, I. J. & PAGEL, M. (2002). The origins of sexual dimorphism in body size in ungulates. *Evolution* **56**, 1276–1285.
- PIENING SCHULER, Y., HAMMER, C., CLAUS, C. & HAMMER, S. (2009). Birth seasonality in captive bovids at the Al Wabra Wildlife Preservation (AWWP), Qatar. In *Proceedings of the International Conference on Diseases of Zoo and Wild Animals*, pp. 1297–1303. Beekse Bergen, The Netherlands.
- POLÁK, J. & FRYNTA, D. (2009). Sexual size dimorphism in domestic goats, sheep, and their wild relatives. *Biological Journal of the Linnean Society* **98**, 872–883.
- POLÁK, J. & FRYNTA, D. (2010). Patterns of sexual size dimorphism in cattle breeds support Rensch's rule. *Evolutionary Ecology* **24**, 1255–1266.
- POST, E. (2003). Timing of reproduction in large mammals. In *Phenology: An Integrative Environmental Science* (ed. M. D. SCHWARTZ), pp. 437–449. Kluwer Academic Publishers, Dordrecht.
- POST, E., BØVING, P. S., PEDERSEN, C. & MACARTHUR, M. A. (2003). Synchrony between caribou calving and plant phenology in depredated and non-depredated populations. *Canadian Journal of Zoology* **81**, 1709–1714.
- POST, E. & FORCHHAMMER, M. C. (2008). Climate changes reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B* **363**, 2369–2375.
- POST, E., PEDERSEN, C., WILMERS, C. C. & FORCHHAMMER, M. C. (2008). Warming, plant phenology and the spatial dimension of trophic mismatch for large herbivores. *Proceedings of the Royal Society B* **275**, 2005–2013.
- POWLESAND, R. G., LLOYD, B. D., BEST, H. A. & MERTON, D. V. (1992). Breeding biology of the Kakapo (*Strigops habroptilus*) on Stewart Island, New Zealand. *Ibis* **135**, 361–373.
- PTAK, G. E., TACCONI, E., CZERNIK, M., TOSCHI, P., MODLINSKI, J. A. & LOI, P. (2012). Embryonic diapause is conserved across mammals. *PLoS ONE* **7**, e33027.
- RACHLOW, J. L. & BOWYER, R. T. (1991). Intraannual variation in timing and synchrony of parturition in Dall's sheep. *Journal of Mammalogy* **72**, 487–492.
- RALLS, K., KRANZ, K. & LUNDRIGAN, B. (1986). Mother-young relationships in captive ungulates: variability and clustering. *Animal Behaviour* **34**, 134–145.
- RANSOM, A. B. (1966). Breeding seasons of white-tailed deer in Manitoba. *Canadian Journal of Zoology* **44**, 59–62.
- RHIND, S. M., ARCHER, Z. A. & ADAM, C. L. (2002). Seasonality of food intake in ruminants: recent developments in understanding. *Nutrition Research Reviews* **15**, 43–65.
- RICE, C. G. (1988). Reproductive biology of Nilgiri tahr (*Hemitragus hylocricus*). *Journal of Zoology* **214**, 269–284.
- ROBBINS, C. T. & ROBBINS, B. L. (1979). Fetal and neonatal growth patterns and maternal reproductive effort in ungulates and subungulates. *American Naturalist* **114**, 101–116.
- ROHLF, F. (2001). Comparative methods for the analysis of continuous variables: geometric interpretations. *Evolution* **55**, 2143–2160.
- RUF, T., FIETZ, J., SCHLUND, W. & BIEBER, C. (2006). High survival in poor years: life history tactics adapted to mast seeding in the edible Dormouse. *Ecology* **87**, 372–381.
- RUTBERG, A. T. (1984). Birth synchrony in American bison (*Bison bison*): response to predation or season? *Journal of Mammalogy* **65**, 418–423.
- RUTBERG, A. T. (1987). Adaptive hypotheses of birth synchrony in ruminants: an interspecific test. *American Naturalist* **130**, 692.
- RYAN, S. J., KNECHTEL, C. U. & GETZ, W. M. (2007). Ecological cues, gestation length, and birth timing in African buffalo (*Syncerus caffer*). *Behavioral Ecology* **18**, 635–644.
- SADLEIR, R. M. F. S. (1969). *The Ecology of Reproduction in Wild and Domestic Mammals*. Methuen & Co, London.
- SAETHER, B.-E., ANDERSEN, R., HJELJORD, O. & HEIM, O. (1996). Ecological correlates of regional variation in life history of the Moose (*Alces alces*). *Ecology* **77**, 1493–1500.
- SANTIAGO-MORENO, J., GÓMEZ-BRUNET, A., TOLEDANO-DÍAZ, A., PICAZO, R., GONZALEZ-BULNES, A. & LÓPEZ-SEBASTIÁN, A. (2006). Seasonal endocrine changes and breeding activity in Mediterranean wild ruminants. *Reproduction in Domestic Animals* **41**, 72–81.
- SHORT, R. V. & HAY, M. F. (1966). Delayed implantation in the roe deer (*Capreolus capreolus*). In *Comparative Biology in Mammals, Symposium of the Zoological Society of London, Vol. 15* (ed. I. W. ROWLANDS), pp. 173–194. Academic Press, London.
- SINCLAIR, A. R., LUDWIG, D. & CLARK, C. W. (2000a). Conservation in the real world. *Science* **289**, 1875.
- SINCLAIR, A. R. E., MDUMA, S. A. & ARCESE, P. (2000b). What determines phenology and synchrony of ungulate breeding in Serengeti? *Ecology* **81**, 2100–2111.
- SKINNER, J. D., AARDE, R. J. V., KNIGHT, M. H. & DOTT, H. M. (1996). Morphometrics and reproduction in a population of Springbok *Antidorcas marsupialis* in the semi-arid southern Kalahari. *African Journal of Ecology* **34**, 312–330.
- SKINNER, J. D., MOSS, D. G. & SKINNER, D. C. (2002). Inherent seasonality in the breeding seasons of African mammals: evidence from captive breeding. *Transactions of the Royal Society of South Africa* **57**, 25.

- SKINNER, D. C., RICHTER, T. A., MALPAUX, B. & SKINNER, J. D. (2001). Annual ovarian cycles in an aseasonal breeder, the springbok (*Antidorcas marsupialis*). *Biology of Reproduction* **64**, 1176–1182.
- SMITH, B. L. & ANDERSON, S. H. (1998). Juvenile survival and population regulation of the Jackson elk herd. *Journal of Wildlife Management* **62**, 1036–1045.
- SOLOUNIAS, N. (2007). Family Giraffidae. In *The Evolution of Artiodactyla* (ed. D. R. PROTHERO), pp. 257–277. John Hopkins University Press, Baltimore.
- SPARKS, T. H. & MENZEL, A. (2002). Observed changes in seasons: an overview. *International Journal of Climatology* **22**, 1715–1725.
- SPINAGE, C. A. (1973). The role of photoperiodism in the seasonal breeding of tropical African ungulates. *Mammal Review* **3**, 71–83.
- TESTA, J. W. (2002). Does predation on neonates inherently select for earlier births? *Journal of Mammalogy* **83**, 699–706.
- TESTA, J. W., BECKER, E. F. & LEE, G. R. (2000). Temporal patterns in the survival of twin and single moose calves (*Alces alces*) in southcentral Alaska. *Journal of Mammalogy* **81**, 162–168.
- THE DUKE OF BEDFORD & MARSHALL, F. H. A. (1942). On the incidence of the breeding season in mammals after transference to a new latitude. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **130**, 396–399.
- UNGERFELD, R., GONZÁLEZ-PENSADO, S., BIELLI, A., VILLAGRÁN, M., OLAZABAL, D. & PÉREZ, W. (2008). Reproductive biology of the Pampas deer (*Ozotoceros bezoarticus*): a review. *Acta Veterinaria Scandinavica* **50**, 16.
- VAN MOURIK, S. A. & STELMASIAK, T. (1990). Endocrine mechanisms and antler cycles in roe deer (*Cervus timorensis*). In *Horns, Pronghorns and Antlers* (eds G. A. BUBENIK and A. B. BUBENIK), pp. 265–297. Springer, New York.
- VERME, L. J. (1965). Reproduction studies on penned White-tailed deer. *Journal of Wildlife Management* **29**, 74–79.
- VERME, L. J. (1969). Reproductive pattern of White-tailed deer related to nutritional plane. *Journal of Wildlife Management* **33**, 881–887.
- VERME, L. J. & OZOGA, J. J. (1987). Relationship of photoperiod to puberty in doe fawn white-tailed deer. *Journal of Mammalogy* **68**, 107–110.
- WECKERLY, F. W. (1998). Sexual-size dimorphism: influence of mass and mating systems in the most dimorphic mammals. *Journal of Mammalogy* **79**, 33–52.
- WESTERN, D. (1979). Size, life history and ecology in mammals. *African Journal of Ecology* **17**, 185–204.
- WILSON, V. J. (2005). *Duikers of Africa (Masters of the African forest floor)*. Zimbi Books, Pretoria.
- ZEDER, M. A. & HESSE, B. (2000). The initial domestication of goats (*Capra hircus*) in the Zagros Mountains 10,000 years ago. *Science* **287**, 2254–2257.
- ZIEBA, D. A., KLOCEK, B., WILLIAMS, G. L., ROMANOWICZ, K., BOLIGLOWA, L. & WOZNIAK, M. (2007). In vitro evidence that leptin suppresses melatonin secretion during long days and stimulates its secretion during short days in seasonal breeding ewes. *Domestic Animal Endocrinology* **33**, 358–365.
- ZUCKERMAN, S. (1952). The breeding seasons of mammals in captivity. *Proceedings of the Zoological Society of London* **122**, 827–950.

VIII. SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article.

Appendix S1. Birth seasonality and biological characteristics of 110 ruminant species.

Appendix S2. Mating season of wild populations.

Appendix S3. Phylogenetic tree used for the phylogenetic generalized least-squares method (PGLS).

Appendix S4. References for Appendices S1–S3.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

(Received 10 April 2012; revised 30 May 2012; accepted 15 June 2012; published online 10 July 2012)

Appendix S1. Birth seasonality and biological characteristics of 110 ruminant species.

Biological characteristics										Seasonality of reproduction				
										Birth windows				
	Body mass (kg) ^a	Sexual size dimorphism ^{ah}	%grass	Mating type ^{ah}	Mother-young relationship ^{ar}	Gestation length (d) ^a	Mid latitude ^a	Seasonality free ranging	Seasonality captivity	BPB50	BPB50 Julian start day	BPB80	BPB80 Julian start day	N
<i>Tragulus javanicus</i>	1.9	0.89 ^{an}	0 ^{d; e}	1 ^{ai}	2	159	-7.4 ^{au}	5 ^{au}	5	180	345	285	125	565
<i>Tragulus napu</i>	5.3	0.98 ^{an}	0 ^{f; e}	1 ^{ai}	2	154	4.2	5 ^{av}	5	175	65	285	45	399
<i>Antilocapra americana</i> *	45 ^b	1.19	15 ^{d; g}	3	2	248	37.2	1 ^{aw; ax}	1	25	150	45	140	1208
<i>Giraffa camelopardalis</i>	965	1.57	0 ^{d; h}	2	2 ^{as}	455	-4.6	4 ^{ay; az}	4	155	120	275	365	3301
<i>Okapia johnstoni</i>	230	0.94 ^{fq}	0 ^d	2 ^{ai}	2	443	-1.6	5 ^{ba}	4	150	150	255	140	428
<i>Moschus moschiferus</i>	13	1.09	4 ⁱ	1	2	176	53.4	1 ^{bb}	1	20	155	50	145	164
<i>Alces alces</i> *	462	1.27	2 ^{d; g}	2	1	235	58.5 ^{fa}	2 (1) ^{bc; bd}	1	20	130	40	125	652
<i>Capreolus capreolus</i> *	22 ^b	1.08	9 ^{d; g}	3	2	298	51.3	2 ^{be}	1	25	140	40	125	457
<i>Hydropotes inermis</i>	13	1.14	50 ^{d; j}	1	2	175	28.9	2 (1) ^{bf}	1	20	145	35	135	767
<i>Rangifer tarandus</i> *	113 ^b	1.45	36 ^{d; g}	3	1 ^{at}	222	64.6	1 ^{bg; bh}	1	25	120	50	115	3527
<i>Odocoileus hemionus</i> *	85	1.36	11 ^{d; g}	2	2	203	41.8	1 ^{bi; bj}	1	20	155	45	145	544
<i>Odocoileus virginianus</i> *	76	1.66	9 ^{d; g}	2	2	201	21.9	2 (1) ^{bk}	1	20	150	55	140	2378
<i>Mazama americana</i>	21	1.00	1 ^d	2	2	225	-9.9	4 ^{bl}	5	175	170	280	210	230
<i>Pudu pudu</i>	9.5	1.00 ^{an}	3 ^k	1 ^j	2	210	-43.7	2 ^{bm}	2	40	120	105	110	715
<i>Ozotoceros bezoarticus</i>	35	1.14 ^{an}	75 ^l	2	2	220	-21.9	4 (3, 2) ^{bn}	3	75	55	160	20	99
<i>Axis axis</i>	70	1.78	70 ^{d; g}	2	2	227	17.3	5 (4, 2) ^{bo; bp}	4	145	355	265	315	5284
<i>Axis porcinus</i>	37	1.34	73 ^m	2	2	220	20.6	2 ^{bq; bp}	4	135	65	270	45	903

<i>Cervus elaphus*</i>	241	1.33	47 ^{d, g}	3	2	236	46.4	2 (1) ^{br, bs; bb}	1	30	150	75	130	4950
<i>Cervus nippon*</i>	53	1.32	50 ^{d, g}	3	2	224	34.0	2 ^{bt; bu}	2	55	145	140	80	3755
<i>Cervus albirostris</i>	161	1.63 ^j	98 ⁿ	3 ^c	2	263	33.7	2 ^{bb}	2	40	160	80	150	221
<i>Elaphurus davidianus</i>	166	1.35	75 ^{d, j}	3	2	286	34.5 ^{fb}	1 ^{bv}	1	25	100	55	90	2045
<i>Cervus eldii</i>	95	1.57	65 ^d	3 ^{ai}	2	240	18.9	2 ^{bw}	2	45	270	95	255	1130
<i>Cervus duvaucelii</i>	169	1.63 ^{an}	75 ^o	3 ^{ai}	2	245	25.8	2 ^{bx}	2	65	140	110	125	1537
<i>Cervus alfredi</i>	46	1.32 ^c	50 ^c	3 ^c	2	243	10.5	2 (1) ^{by}	4	130	250	245	165	160
<i>Cervus unicorn</i>	176	1.32	28 ^{m, p}	3	2	246	12.0	2 ^{bp}	4	105	280	230	205	560
<i>Cervus timorensis</i>	66	1.38	75 ^c	3	2	250	-7.4	3 ^{bz}	3	65	255	225	155	195
<i>Dama dama*</i>	55 ^b	1.43	46 ^{d, g}	3	2	230	48.3	2 (1) ^{ca; cb}	1	20	160	45	155	4459
<i>Dama mesopotamica*</i>	74	1.43 ^c	46 ^c	3 ^c	2	225 ^{fc}	33.6 ^{fm}	2 ^{cc}	2	35	125	70	120	241
<i>Muntiacus muntjak</i>	18	1.13 ^{an}	10 ^d	1	2	210	11.8	5 ^{bp}	5	170	140	280	275	368
<i>Muntiacus reevesi</i>	14	0.80	10 ^{f, g}	1 ^{ai}	2	214	27.2	4 ^{od}	4	135	115	265	95	3294
<i>Elaphodus cephalophus</i>	23	1.25 ^{fv}	10 ^q	1 ^{ai}	2	181	27.9	2 ^{ce}	3	90	105	205	95	203
<i>Cephalophus dorsalis</i>	20	0.96 ^{ak}	0 ^r	1 ^{ak}	2	238	1.1	5 (4) ^{ak; cg}	5	150	105	275	105	174
<i>Cephalophus silvicultor</i>	62	0.73 ^{fs}	1 ^{s, r}	1 ^{ai}	2	234	-1.9	5 (4) ^{al}	5	125	290	255	290	214
<i>Cephalophus maxwellii</i>	8.4	0.94 ^{ak}	1 ^r	1 ^{ak}	2	205	9.1	4 ^{ch}	5	145	365	270	240	257
<i>Cephalophus monticola</i>	4.9	0.87 ^{an}	0 ^r	1 ^l	2	205	-12.5	5 (4) ^{ak; cg}	5	155	100	265	75	316
<i>Bos frontalis</i>	800 ^b	1.49	66 ^t	2	1	274	15.4	4 ^{ad; cj; ck}	4	145	110	265	70	908
<i>Bos javanicus</i>	667 ^b	1.67 ^{fs}	80 ^d	2 ^{ai}	1	297	8.2	2 ^{cl}	4	145	40	250	350	885
<i>Bos grunniens*</i>	500	2.11 ^{fw}	86 ⁿ	2 ^{ai}	1	274	34.3	2 ^{cm}	3	85	115	160	90	1818
<i>Bison bison*</i>	625	1.65	84 ^{d, g}	2	1 ^{as}	283	44.3	2 ^{cn}	2	45	115	115	105	4474
<i>Bison bonasus*</i>	676	1.70 ^{fs}	68 ^{d, g}	2 ^{ai}	1	266	51.1 ^{fn}	2 ^{co}	2	70	130	150	120	1694
<i>Bubalus bubalis</i>	930	1.50 ^{ft}	80 ^u	2 ^{ai}	1	320	21.0	2 ^{cp}	4	110	185	235	100	549
<i>Bubalus depressicornis</i>	257	1.08 ^{ft}	4 ^v	2 ^c	1	296	-1.3	5 ^{cq}	5	155	190	270	70	277
<i>Syncerus caffer</i>	593	1.30	90 ^{d, g}	2	1	337	-8.8	4 (3, 2) ^{cr; bh}	5	175	185	290	185	937
<i>Boselaphus tragocamelus</i>	182	1.44	29 ^{v; d; h}	3	2	248	22.4	2 ^{ad}	3	105	195	240	90	2971
<i>Tetracerus quadricornis</i>	19 ^b	1.06 ^{fs}	9 ^v	1 ^{ai}	1	207	22.3	4 ^{cs}	5	100	340	220	350	175
<i>Taurotragus derbianus</i>	646	1.55 ^{an}	5 ^r	2 ^c	2	275 ^{td}	8.8	3 ^{ct}	3	85	180	170	130	164
<i>Taurotragus oryx</i>	563	1.47	10 ^{fj; fk; fl}	2	2	271	-10.1	4 ^{cu; cr}	4	115	55	235	5	3546
<i>Tragelaphus strepsiceros</i>	206	1.51	5 ^{d; g}	2	2	271 ^{fr}	-7.0	4 (3, 2) ^{cv; ow}	3	70	195	175	180	2510
<i>Tragelaphus imberbis</i>	94	1.14	10 ^{d; h}	2	2	222	3.9	5 ^{cx}	4	140	255	270	255	826
<i>Tragelaphus eurycerus</i>	271	1.25	20 ^w	2	2 ^{as}	285	2.1	5 ^{cy}	4	140	5	250	340	1595
<i>Tragelaphus spekii</i>	76	1.89	67.5 ^d	2	2	241	-3.3	5 ^{ay}	4	160	365	275	320	3165
<i>Tragelaphus scriptus</i>	43	1.58	10 ^u	2	2	183	-9.4	5 ^{cz; ay}	5	165	5	270	10	187

<i>Tragelaphus angasii</i>	88	1.83 ^{an}	20 ^d	2	2	220	-22.0	5 (4) ^{da; db}	170	15	275	295	2057
<i>Kobus ellipsiprymnus</i>	202	1.33	80 ^{d; h}	3	2	270	-6.5	4 ^{cr; bh}	150	145	275	20	2841
<i>Kobus leche</i>	89	1.31	95 ^r	3	2	230	-14.2	4 (2) ^{dc; db; ay}	150	95	270	10	2139
<i>Kobus megaceros</i>	86	1.41 ^{ft}	95 ^r	3 ^{ai}	2	237 ^{fe}	8.0	2 ^{dd}	135	295	235	275	1527
<i>Kobus kob</i>	80	1.47	95 ^d	3	2	255	6.2	5 ^{de; df}	170	95	280	45	450
<i>Redunca fulvorufula</i>	30 ^b	1.03	99 ^{d; h}	2	2	223	-12.3	5 (4) ^{d; cw}	90	165	205	100	220
<i>Aepyceros melampus</i>	53	1.33	60 ^{d; g}	3	2	197	-12.9	4 ^{dh; da; di; cr}	55	135	190	120	2648
<i>Antidorcas marsupialis</i>	34	1.21	30 ^d	3	2	170	-20.7	4 ^{dj}	85	110	235	95	1856
<i>Antilope cervicapra</i>	38 ^b	1.09 ^{an}	61 ^{v; d; g}	3 ^{ai}	2	167	18.8	2 ^{dk}	145	45	265	65	7042
<i>Gazella dorcas</i>	16	1.31 ^{ci}	25 ^r	3 ^{ai}	2 ^{at}	172	23.6	2 ^{di; dm}	160	40	275	45	815
<i>Gazella spekei</i>	20 ^b	1.28 ^{fs}	50 ^r	3 ^{ai}	2	179	6.5		145	95	270	80	309
<i>Gazella arabica</i>	24	1.31 ^c	50 ^c	3 ^{ai}	2	170 ^c	13.7		135	125	245	120	406
<i>Gazella subgutturosa</i>	27	1.18 ^{cf}	50 ^f	3 ^{ai}	2	160	32.1	2 (1) ^{dn}	25	120	75	90	1807
<i>Gazella cuvieri</i>	20 ^c	1.44 ^{ft}	50 ^c	3 ^c	2	170 ^{fr}	30.4		105	45	230	15	529
<i>Gazella leptoceros</i>	24 ^b	1.30 ^{ft}	50 ^r	3 ^{ai}	2	167	24.8		50	75	200	65	1127
<i>Gazella thomsonii</i>	23	1.20	86 ^{w; h}	3	2	166	1.9	4 ^{bh}	115	100	250	90	1960
<i>Nanger dama</i>	73 ^b	1.33 ^{fs}	48 ^r	3 ^{ai}	2	195	20.2	4 (3, 2, 1) ^{do}	130	170	250	70	2747
<i>Nanger soemmerringii</i>	42	1.35 ^{fs}	50 ^r	3 ^c	2	193	9.8		95	190	220	115	265
<i>Nanger granti</i>	56	1.45	50 ^{d; g}	3	2	191	0.3	5 (3) ^{bh; dp; cx}	125	140	250	140	1031
<i>Litocranius walleri</i>	39	1.45	0 ^{d; h}	3	2	204	3.0	5 ^{dq}	150	145	265	125	552
<i>Saiga tatarica</i> *	38 ^b	1.39	26 ^y	3	2	148	46.5	1 ^{dr}	15	130	30	120	414
<i>Madoqua guentheri</i>	4.6	0.82 ^{am}	5 ^u	1 ^{am}	2	175	4.3	4 (2) ^{am; dp}	170	205	260	95	266
<i>Madoqua kirkii</i>	4.9 ^b	0.93	17 ^r	1	2 ^{as}	171	-6.6	4 (2) ^{ds; dp}	165	285	255	255	652
<i>Neotragus moschatus</i>	5.6	0.93 ^{an}	0 ^d	1 ^{an}	2	180	-13.3	4 ^{dp}	170	45	275	305	292
<i>Oreotragus oreotragus</i>	14	1.00	5 ^u	1	2	214	-7.8	4 ^{dp}	150	180	270	120	212
<i>Addax nasomaculatus</i>	96	1.50 ^{aj}	80 ^r	3 ^{ai}	2	263	20.5	4 ^{dt; du}	135	360	245	340	3080
<i>Oryx gazelle</i>	195 ^b	1.10	82 ^{d; h}	3	2	274	-8.3	4 ^{cz}	155	35	265	360	2748
<i>Oryx leucoryx</i>	76	1.06 ^{an}	75 ^r	3 ^{ai}	2	240	20.6 ^{fo}	4 ^{dv}	130	20	240	330	1340
<i>Oryx dammah</i>	200	1.10 ^{fs}	75 ^r	3 ^{ai}	2 ^{as}	248	16.2	4 ^{dw}	125	85	240	365	3957
<i>Hippotragus niger</i>	236	1.08	93 ^d	3	2	273	-13.6	2 ^{dx; ay; dy}	155	105	270	25	2555
<i>Hippotragus equinus</i>	264	1.02	85 ^r	3	2	278	-4.6	5 (4) ^{dy; ay}	155	50	270	340	1031
<i>Alcelaphus buselaphus</i>	161	1.07	97 ^{d; h}	3	2	243	5.8	4 ^{cr; bh}	90	105	195	95	365
<i>Damaliscus lunatus</i>	78	1.18	99.3 ^u	3	1	238	-16.5	2 ^{d; bh}	65	115	165	100	166
<i>Damaliscus pygargus</i> *	85 ^b	1.11 ^{an}	100 ^d	3 ^{ai}	1	241	-29.5	2 ^{eat; eb; ec}	65	130	155	120	1593
<i>Connochaetes taurinus</i>	199	1.35	90 ^{d; g}	3 ^{ai}	1 ^{at}	251	-15.6	1 (2) ^{ed; cr; ay}	50	155	125	130	1652

<i>Connochaetes gnou</i> *	157	1.29 ^{an}	81 ^r	3 ^{ai}	1	257	-28.8	1 ^{ee}	2	50	165	110	120	471
<i>Budorcas taxicolor</i> *	295	1.18 ^{ao}	30 ^c	3 ^{ao}	1	236	30.2	2 ^{ef, bb}	2	60	35	135	15	271
<i>Ovis orientalis musimon</i> *	50 ^c	1.51 ^{fu}	69 ^u	2	1	153 ^{fg}	41.2 ^{fp}	1 ^{eg, eh}	2	45	75	90	55	3351
<i>Ovis orientalis gmelini</i>	50 ^b	1.59 ^{fu}	69 ^d	2	1	158 ^{fh}	36.1 ^{fp}	1 ^{ei}	1	15	130	30	125	769
<i>Ovis orientalis vignei</i> *	52	1.54 ^{fu}	70 ^u	2	1	165	34.2 ^{fp}	1 ^{ej}	1	20	105	40	100	748
<i>Ovis canadensis</i> *	75	1.43	67 ^g	2	1	177	40.5	1 ^{ek}	1	25	125	55	105	1239
<i>Ovis dalli</i> *	70	1.57	56 ^{w, g}	2	1	172	60.6	1 ^{el, em}	1	30	125	85	95	1161
<i>Capra aegagrus</i> *	47	1.70	28 ^u	2	1	156	33.6	2 ^{en}	3	45	100	150	55	1499
<i>Capra ibex</i> *	72 ^b	2.11 ^{an}	60 ^{w, g}	2 ^{ai}	1 ^{at}	168	45.8	1 ^{eo}	1	20	140	45	125	1517
<i>Capra sibirica</i> *	130	2.04 ^x	50 ^z	2 ^{ai}	1	175	43.5	1 ^{ep, eq}	1	20	135	45	125	986
<i>Capra caucasica</i> *	58	1.57 ^{ft}	80 ^{aa}	2 ^{ai}	1	155	42.3	2 ^{er}	1	20	135	40	125	858
<i>Capra cylindricornis</i> *	58	1.66 ^{an}	47.5 ^{ab}	2 ^{ab}	1	155	42.1 ^{es}	2 ^{es}	1	20	145	40	135	246
<i>Capra nubiana</i>	48	2.36 ^{fu}	60 ^c	2 ^{ai}	1	150	23.5	2 ^{(1)et}	1	30	85	55	75	1095
<i>Capra falconeri</i>	54	2.27 ^{an}	60 ^c	2 ^{ai}	1	168	33.9	2 ^{(1)eu}	1	20	140	40	130	2458
<i>Hemitragus hylocrius</i>	74	2.00 ^{fs}	64 ^{ac}	2 ^{ai}	1	186	10.0	2 ^{ev}	5	150	135	260	50	178
<i>Hemitragus jemlahicus</i>	68	2.03	75 ^{d, ad}	2	1 ^{at}	196	30.1	2 ^{bb}	2	30	150	55	140	1436
<i>Pseudois nayaur</i>	52	1.35	68 ^{ae}	2	1	160	34.4	1 ^{ae}	1	30	165	60	150	259
<i>Ammotragus levia</i>	94	1.46	42 ^{af}	2	1	156	24.8	1 ^{(2)ew}	2	55	70	145	30	4804
<i>Rupicapra rupicapra</i>	33	1.21	74 ^{d, g}	3 ^{ai}	1	178	43.5	1 ^{ex, ey}	1	25	130	55	110	449
<i>Oreamnos americanus</i> *	72	1.39	61 ^g	2	1	181	54.1	1 ^{ez}	1	15	145	35	135	714
<i>Naemorhedus caudatus</i>	27	1.11 ^{an}	50 ^c	2 ^c	1	180 ^{fi}	42.0	2 ^{bb}	2	45	135	90	125	264
<i>Capricornis crispus</i>	43	1.00	3.1 ^{ag}	2 ^{ap}	1	213	36.8	2 ^{ap}	2	30	125	90	125	97
<i>Ovibos moschatus</i> *	313	1.47	62 ^g	3 ^{aq}	1	256	71.7	2 ^{aq}	2	30	135	70	125	637

References (see online Appendix S4):

- ^a Jones *et al.* (2009); ^b Ernest (2003); ^c estimated from close relative as sufficient data were not available; ^d Hofmann *et al.* (2008); ^e Nordin (1978); ^f Clauss *et al.* (2008); ^g Van Wieren (1996); ^h Owen-Smith (1997); ⁱ Green (1987); ^j Geist (1998); ^k Eldridge *et al.* (1987); ^l Jackson & Giulietti (1988); ^m Dinerstein (1982); ⁿ Harris & Miller (1995); ^o Wegge, Shrestha & Moe (2006); ^p Padmalal, Takatsuki & Jayasekara (2003); ^q Sheng & Lu (1982); ^r Gagnon & Chew (2000); ^s Gautier-Hion, Emmons & Dubost (1980); ^t Chetri (2006); ^u Clauss *et al.* (2010); ^v Pujaningsih *et al.* (2009); ^w Solanki & Naik (1998); ^x Fedosenko & Blank (2001); ^y Bannikov *et al.* (1967); ^z estimated from Fedosenko & Blank (2005); ^{aa} Weinberg (2004); ^{ab} Weinberg (2008); ^{ac} Sumithran (1997); ^{ad} Schaller (1967); ^{ae} Wang & Hoffmann (1987); ^{af} Ogren (1962); ^{ag} Ochiai (1999); ^{ah} Weckerly (1998); ^{ai} Grzimek (1979/1980); ^{aj} Krausman & Casey (2007); ^{ak} Wilson (2005); ^{al} Lumpkin & Kranz (1984); ^{am} Kingswood & Kumamoto (1996); ^{an} Loison *et al.* (1999); ^{ao} Neas & Hoffmann (1987); ^{ap} Jass & Mead (2004); ^{aq} Lent (1988); ^{ar} Stoner, Caro & Graham (2003); ^{as} Ralls, Kranz & Lundrigan (1986); ^{at} Ralls, Lundrigan & Kranz (1987); ^{au} Duckworth *et al.* (2008); ^{av} Lutz (2001); ^{aw} Kitchen (1974); ^{ax} Buechner (1950); ^{ay} Ansell (1960); ^{az} Fairall (1968); ^{ba} Bodmer & Rabb (1992); ^{bb} Unknown Author (2005); ^{bc} Haagenrud & Markgren (1974); ^{bd} Stringham (1974); ^{be} Linnell & Andersen (1998); ^{bf} Kirkwood (2007); ^{bg} Lent (1966);

^{bh} Sinclair, Mduma & Arcese (2000); ^{bi} Jordan (1976); ^{bj} Anderson & Olof (1984); ^{bk} Illige (1951); ^{bl} Gardner (1971); ^{bm} Pollard (1999); ^{bn} Ungerfeld *et al.* (2008); ^{bo} Loudon (1988); ^{bp} Mishra & Wemmer (1987); ^{bq} Dhungel & O'Gara (1991); ^{br} Guinness, Clutton-Brock & Albon (1978); ^{bs} Clutton-Brock, Guinness & Albon (1983); ^{bt} Evtushevsky (1974); ^{bu} Maruyama *et al.* (1975); ^{bv} Wemmer *et al.* (1989); ^{bw} Worrel (2004); ^{bx} Ferraino & Yahnke (2007); ^{by} Key (2003); ^{bz} Reyes (2002); ^{ca} Chapman & Chapman (1975); ^{cb} Sterba & Klusak (1984); ^{cc} Perelberg *et al.* (2003); ^{cd} Pei *et al.* (1995); ^{ce} Lundrigan & Oas (2003); ^{cf} Kingswood & Blank (1996); ^{cg} Kranz & Lumpkin (1982); ^{ch} Aeschlimann (1963); ^{ci} Tom-Tov (1995); ^{cj} Vairavel (1998); ^{ck} National Research Council (1983); ^{cl} Choquent (1993); ^{cm} Zi (2003); ^{cn} Rutberg (1984); ^{co} Balčiauskas (1999); ^{cp} Singh, Singh & Bains (1985); ^{cq} Miller (2002); ^{cr} Spinage (1973); ^{cs} Leslie & Sharma (2009); ^{ct} Altan (2000); ^{cu} Pappas (2002); ^{cv} Simpson (1968); ^{cw} Kingdon (1982); ^{cx} Leuthold & Leuthold (1975); ^{cy} Turkalo & Turkalo (1999); ^{cz} Skinner, Moss & Skinner (2002); ^{da} Anderson (1979); ^{db} Leuthold (1977); ^{dc} Sayer & Lavieren (1975); ^{dd} Corrie & Yahnke (2004); ^{de} Modha & Eltringham (1976); ^{df} Leuthold (1967); ^{dg} Irby (1979); ^{dh} Murray (1982); ^{di} Moe, Rutina & Du Toit (2007); ^{dj} Skinner *et al.* (1996); ^{dk} Prakash (1960); ^{dl} Mendelssohn & Yom-Tov (1987); ^{dm} Yom-Tov, Mendelssohn & Groves (1995); ^{dn} Kingswood & Blank (1996); ^{do} Cano & Cano (2007); ^{dp} Spinage (1986); ^{dq} Penfold (2005); ^{dr} Sokolov (1974); ^{ds} Kingswood & Kumamoto (1997); ^{dt} Manski (1991); ^{du} Asa (1996); ^{dv} Vié (1996); ^{dw} Morrow (1999); ^{dx} Sekulic (1978); ^{dy} Wilson & Hirst (1977); ^{dz} Child, Robbel & Hepburn (1972); ^{ea} Du Plessis (1972); ^{eb} David (1973); ^{ec} David (1975); ^{ed} Estes (1976); ^{ee} Richter (1971); ^{ef} Wang *et al.* (2005); ^{eg} Langbein, Streich & Scheibe (1998); ^{eh} Langbein (1998); ^{ei} Valdez, Cardenas & Sanchez (1991); ^{ej} Awan & Festa-Bianchet (2006); ^{ek} Shackleton (1985); ^{el} Bowyer & Leslie (1992); ^{em} Bunnell (1980); ^{en} Edge (1990); ^{eo} Stüwe & Grodinsky (1987); ^{ep} Dmitriev (1938); ^{eq} Roberts (1977); ^{er} Adler & Bar-Oz (2009); ^{es} Weinberg (2002); ^{et} Habibi (1997); ^{eu} Coblenz (1980); ^{ev} Rice (1988); ^{ew} Gray & Simpson (1980); ^{ex} Hardenberg (2000); ^{ey} Boschi (2003); ^{ez} Rideout & Hoffmann *et al.* (2008); ^{fb} Zhigang & Harris (2008); ^{fc} Cavendish (2001); ^{fd} Angwafo (2006); ^{fe} Falchetti & Mostacci (1993); ^{ff} Furley (1986); ^{fg} Reavill (2000); ^{fh} Flores-Foxworth *et al.* (1995); ^{fi} Mead (1989); ^{fj} Codron *et al.* (2007); ^{fk} Watson & Owen-Smith (2002); ^{fl} Sponheimer *et al.* (2003); ^{fm} Rabiei (2010); ^{fn} Olech (2008); ^{fo} IUCN SSC Antelope Specialist Group (2008); ^{fp} Valdez (2008); ^{fq} ISIS data were used; ^{fr} Dittrich (1972); ^{fs} Bro-Jørgensen (2008); ^{ft} Polák & Frynta (2009); ^{fv} Carranza (1996); ^{fw} Leslie & Schaller (2009).

Body mass: median of the mean male and female body mass; sexual size dimorphism: ratio between body mass of males and females; % grass: percentage of grass in the natural diet of a species; Mating type: 1 = monogamous, 2 = males follow more than one female but no harem or territory is defended, 3 = harem or territory is defended; Mother-young relationship: 1 = follower, 2 = hider; Mid latitude: centre of the latitude range that a species inhabits; Seasonality of reproduction (categories for free ranging and captive) from 1: highly seasonal to 5: aseasonal species (for details see text and Fig. 2; note that for comparisons between free-ranging and captive populations, the reported category not in parentheses was used, which were considered most typical for the species; categories in parentheses represent other patterns described for the species in the literature); BPB: birth peak breadth, measure of the degree of reproductive seasonality as number of days (measured as consecutive 5-day blocks) which included the required percentage of birth (50% or 80%); Julian start day: first Julian day (of the year) of the respective BPB; N: number of observed birth events in the captive population. For species that are marked with an asterisk (*) data on the exact start of mating season and the latitude of origin of at least one free-ranging population were available (Appendix S2). These data were used for further analyses.

Appendix S2. Mating season of wild populations

Start of mating seasons for populations of seasonal ruminants for which the exact start date of the mating season (sometimes calculated as first observed birth event minus the average gestation length) and the exact mid latitude of the origin of the population were available.


Species	Start of mating season (Julian day)	Latitude of origin	Reference
<i>Antilocapra americana</i>	267	40.1	Fairbanks (1993)
<i>Alces alces</i>	267	63.8	Bowyer, Van Ballenberghe & Kie (1998)
<i>Capreolus capreolus</i>	289	63.7	Linnell & Andersen (1998)
<i>Rangifer tarandus</i>	269	62.2	Reimers, Klein & Sørungård (1983)
<i>Odocoileus hemionus</i>	329	32.9	Bowyer (1991)
<i>Odocoileus virginianus</i>	300	47.1	Carstensen Powell & DelGiudice, 2005)
<i>Cervus elaphus</i>	256	57	Guinness <i>et al.</i> (1978)
<i>Cervus nippon</i>	252	35	Asada & Ochiai (1996)
<i>Dama dama</i>	280	37	Braza, San Jose & Blom (1988)
<i>Dama mesopotamica</i>	225	33	Perelberg <i>et al.</i> (2003)
<i>Bos grunniens</i>	183	32.5	Zi (2003)
<i>Bison bison</i>	193	47.3	Rutberg (1984)
<i>Bison bonasus</i> *	282	56.1	Balčiauskas (1999)
<i>Saiga tatarica</i>	339	48	Bekenov, Grachev & Milner-Gulland (1998)
<i>Damaliscus pygargus</i>	260 ^a	-25.9	Du Plessis (1972)
<i>Connochaetes gnou</i>	246 ^a	-30.5	Vrahimis & Kok (1994)
<i>Budorcas taxicolor</i>	160	33.7	Wang <i>et al.</i> (2005)
<i>Ovis orientalis musimon</i>	277	40.4	Santiago-Moreno <i>et al.</i> (2001)
<i>Ovis orientalis vignei</i> *	231	25	Awan & Festa-Bianchet (2006)
<i>Ovis canadensis</i> *	332	58.8	Bunnell (1982)
<i>Ovis dalli</i> *	323	64	Rachlow & Bowyer (1991)
<i>Capra aegagrus</i>	224	25.6	Edge (1990)
<i>Capra ibex</i>	336	46.5	Giocometti & Ratti (1994)
<i>Capra sibirica</i>	305	45	Fedosenko & Blank (2001)
<i>Capra caucasica</i>	363	43.4	Dinnik (1887)
<i>Capra cylindricornis</i>	369	42.8	Weinberg (2002)
<i>Oreamnos americanus</i>	324	54	Côté & Festa-Bianchet (2001)
<i>Ovibos moschatus</i> *	205	60.1	Lent (1988)

Species marked with an asterisk (*) are outliers. Statistical analyses were performed with and without these species.

^acorrected for transformation to the northern hemisphere.

See Appendix S4 for references.

Appendix S3. Phylogenetic tree used for the phylogenetic generalized least-squares method (PGLS).



Appendix S4. References for Appendices S1-S3.

- ADLER, D. S. & BAR-OZ, G. (2009). Seasonal patterns of prey acquisition and inter-group competition during the middle and upper Palaeolithic of the Southern Caucasus. In *The Evolution of Hominin Diets. Vertebrate Paleobiology and Paleoanthropology* (ed. J.-J. Hublin and M. P. Richards), pp. 127-140. Springer Netherlands.
- AESCHLIMANN, A. (1963). Observations sur *Philantomba maxwelli* (Hamilton-Smith) une antilope de la forêt éburnéenne. *Acta Tropica* **20**, 341-368.
- ALTAN, B. (2000). *Taurotragus derbianus*, pp. http://animaldiversity.ummz.umich.edu/site/accounts/information/Taurotragus_derbianus.html. Animal Diversity Web.
- ANDERSON, A. E. & OLOF, C. W. (1984). *Odocoileus hemionus*. *Mammalian Species*, 1-9.
- ANDERSON, J. L. (1979). Reproductive seasonality of the Nyala *Tragelaphus angasi*; The interaction of light, vegetation phenology, feeding style and reproductive physiology. *Mammal Review* **9**, 33-46.
- ANGWAFO, T. E. (2006). Status of wildlife and its utilisation in Faro and Benoué National Parks north Cameroon: case study of the Derby eland (*Taurotragus derbianus gigas* Gray, 1947) and the African wild dog (*Lycaon pictus* Temminck, 1840), Brandenburg Technical University.
- ANSELL, W. F. H. (1960). The breeding of some larger mammals in northern Rhodesia. *Proceedings of the Zoological Society of London* **134**, 251-274.
- ASA, C. S. (1996). Ovulatory cycles and anovulatory periods in the addax (*Addax nasomaculatus*). *Journal of Reproduction and Fertility* **107**, 119.
- ASADA, M. & OCHIAI, K. (1996). Conception dates of sika deer on the Boso Peninsula, central Japan. *Mammal Study* **21**, 153-159.
- AWAN, G. A. & FESTA-BIANCHET, M. (2006). Lambing seasons in subspecies of Urial (*Ovis vignei*) in Pakistan. *Journal Mountain Ecology* **8**, 27-32.
- BALČIAUSKAS, L. (1999). European bison (*Bison bonasus*) in Lithuania: status and possibilities of range extension. *Acta Zoologica Lituanica* **9**, 3-18.
- BANNIKOV, A., ZHIRNOV, L. V., LEBEDEVA, L. S. & FANDEEV, A. A. (1967). Biology of the Saiga. *Jerusalem: Israel Program for Scientific Translations*.
- BEKENOV, A. B., GRACHEV, I. A. & MILNER-GULLAND, E. J. (1998). The ecology and management of the Saiga antelope in Kazakhstan. *Mammal Review* **28**, 1-52.
- BODMER, R. E. & RABB, G. B. (1992). *Okapia johnstoni*. *Mammalian Species*, 1-8.
- BOSCHI, C. (2003). The spatial patterns of Alpine chamois (*Rupicapra rupicapra rupicapra*) and their influence on population dynamics in the Swiss National Park. *Mammalian biology* **68**, 16.
- BOWYER, R. T. (1991). Timing of parturition and lactation in southern mule deer. *Journal of Mammalogy* **72**, 138-145.
- BOWYER, R. T. & LESLIE, D. M., JR. (1992). *Ovis dalli*. *Mammalian Species*, 1-7.
- BOWYER, R. T., VAN BALLEMBERGHE, V. & KIE, J. G. (1998). Timing and synchrony of parturition in Alaskan moose: long-term versus proximal effects of climate. *Journal of Mammalogy* **79**, 1332-1344.
- BRAZA, F., SAN JOSE, C. & BLOM, A. (1988). Birth measurements, parturition dates, and progeny sex ratio of *Dama dama* in Doñana, Spain. *Journal of Mammalogy* **69**, 607-610.
- BRO-JØRGENSEN, J. (2007). The intensity of sexual selection predicts weapon size in male bovids. *Evolution* **61**, 1316-1326.
- BRO-JØRGENSEN, J. (2008). Dense habitats selecting for small body size: a comparative study on bovids. *Oikos* **117**, 729-737.

- BUECHNER, H. K. (1950). Life history, ecology, and range use of the pronghorn antelope in Trans-Pecos Texas. *The American Midland Naturalist* **43**, 257.
- BUNNELL, F. L. (1980). Factors controlling lambing period of Dall's sheep. *Canadian Journal of Zoology* **58**, 1027-1031.
- BUNNELL, F. L. (1982). The lambing period of Mountain sheep: synthesis, hypotheses, and tests. *Canadian Journal of Zoology* **60**, 1-14.
- CANO, M. & CANO. (2007). Establishment of a group of dama gazelles for reintroduction in Senegal. *International Zoo Yearbook* **32**, 98.
- CARRANZA, J. (1996). Sexual selection for male body mass and the evolution of litter size in mammals. *The American Naturalist* **148**, 81-100.
- CARSTENSEN POWELL, M. & DELGIUDICE, G. D. (2005). Birth, morphologic, and blood characteristics of free-ranging white-tailed deer neonates. *Journal of Wildlife Diseases* **41**, 171-183.
- CAVENDISH, M. (2001). *Endangered wildlife and plants of the world*. Marshall Cavendish Corporation, Tarrytown, NY.
- CHAPMAN, D. I. & CHAPMAN, N. G. (1975). *Fallow Deer, their history, distribution and biology*. Terence Dalton Ltd., Lavenham, Suffolk.
- CHETRI, M. (2006). Diet analysis of Gaur, *Bos gaurus gaurus* (Smith, 1827) by micro-histological analysis of fecal samples in Parsa Wildlife Reserve, Nepal. *Our Nature* **4**, 20-28.
- CHILD, G., ROBBEL, H. & HEPBURN, C. P. (1972). Observations on the biology of the Tsessebe, *Damaliscus lunatus lunatus*, in northern Botswana. *Mammalia* **36**, 342-388.
- CHOQUENT, D. (1993). Growth, body condition and demography of wild Banteng (*Bos javanicus*) on Cobourg Peninsula, northern Australia. *Journal of Zoology* **231**, 533-542.
- CLAUSS, M., HOFMANN, R., STREICH, W., FICKEL, J. & HUMMEL, J. (2008). Higher masseter muscle mass in grazing than in browsing ruminants. *Oecologia* **157**, 377-385.
- CLAUSS, M., MÜLLER, K., FICKEL, J., STREICH, W. J., HATT, J. M. & SÜDEKUM, K. H. (2010). Macroecology of the host determines microecology of endobionts: protozoal faunas vary with wild ruminant feeding type and body mass. *Journal of Zoology*, no-no.
- CLUTTON-BROCK, T. H., GUINNESS, F. E. & ALBON, S. D. (1983). The costs of reproduction to Red deer hinds. *Journal of Animal Ecology* **52**, 367-383.
- COBLENTZ, B. E. (1980). A unique ungulate breeding pattern. *The Journal of Wildlife Management* **44**, 929-933.
- CODRON, D., CODRON, J., LEE-THORP, J. A., SPONHEIMER, M., DE RUITER, D., SEALY, J., GRANT, R. & FOURIE, N. (2007). Diets of savanna ungulates from stable carbon isotope composition of faeces. *Journal of Zoology* **273**, 21-29.
- CORRIE, J. & YAHNKE, C. (2004). *Kobus megaceros*. Animal Diversity Web.
- CÔTÉ, S. D. & FESTA-BIANCHET, M. (2001). Birthdate, mass and survival in mountain goat kids: effects of maternal characteristics and forage quality. *Oecologia* **127**, 230-238.
- DAVID, J. H. M. (1973). The Behaviour of the Bontebok, *Damaliscus dorcas dorcas*, (Pallas 1766), with Special Reference to Territorial Behaviour*. *Zeitschrift für Tierpsychologie* **33**, 38-107.
- DAVID, J. H. M. (1975). Observations on mating behaviour, parturition, suckling and the mother-young bond in the Bontebok (*Damaliscus dorcas dorcas*). *Journal of Zoology* **177**, 203-223.
- DHUNGEL, S. K. & O'GARA, B. W. (1991). Ecology of the Hog deer in Royal Chitwan National Park, Nepal. *Wildlife Monographs*, 3-40.
- DINERSTEIN, E. (1982). Deer, plant phenology and succession in the lowland forests of Nepal. In *Biology and management of the Cervidae* (ed. C. Wemmer), pp. 272- 288. Smithsonian Institution Press, Washington D.C.

- DINNIK, H. (1887). On the Caucasian mountain-goat (*Capra caucasica*). *Journal of Natural Historie Series* **5** **19**, 450-461.
- DITTRICH, L. (1972). Gestation periods and age of sexual maturity of some African antelopes. *International Zoo Yearbook* **12**, 184-187.
- DMITRIEV, V. V. (1938). The ungulates of the Altai Reserve and adjoining areas. In *Proceedings of the Altai Reserve*, vol. 1, pp. 171-259.
- DU PLESSIS, S. S. (1972). Ecology of Blesbok with special reference to productivity. *Wildlife Monographs* **30**, 1-70.
- DUCKWORTH, J. W., HEDGES, S., TIMMINS, R. J. & SEMIADI, G. (2008). *Tragulus javanicus*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.
- EDGE, W. D. (1990). Population characteristics and group composition of *Capra aegagrus* in Kirthar National Park, Pakistan. *Journal of Mammalogy* **71**, 156.
- ELDRIDGE, W. D., MACNAMARA, M. M. & PACHECO, N. V. (1987). Activity patterns and habitat utilization of Pudus (*Pudu puda*) in south-central Chile. In *Biology and management of the cervidae* (ed. C. M. Wemmer), pp. 352-369. Smithsonian Inst. Press., Washington, D.C.
- ERNEST, S. K. M. (2003). Life history characteristics of placental nonvolant mammals. *Ecology* **84**, 3402.
- ESTES, R. D. (1976). The significance of breeding synchrony in the Wildebeest. *African Journal of Ecology* **14**, 135-152.
- EVTUSHEVSKY, N. N. (1974). Reproduction of *Cervus nippon hortulorum* SW. under conditions of the middle Dnieper area. *Vestnik Zoologii* **7**, 23-28.
- FAIRALL, N. (1968). The reproductive season of some mammals in the Kruger National Park. *Zoologica Africana* **3**, 189-210.
- FAIRBANKS, W. S. (1993). Birthdate, birthweight, and survival in pronghorn fawns. *Journal of Mammalogy* **74**, 129-135.
- FALCHETTI, E. & MOSTACCI, B. (1993). The Nile lechwe *Kobus megaceros*: PVA factors and guidelines to captive management. *International Zoo Yearbook* **32**, 60-69.
- FEDOSENKO, A. K. & BLANK, D. A. (2001). *Capra sibirica*. *Mammalian Species* **675**, 1-13.
- FEDOSENKO, A. K. & BLANK, D. A. (2005). *Ovis ammon*. *Mammalian Species*, 1-15.
- FERRAINO, A. & YAHNKE, C. (2007). *Rucervus duvaucelii*. Animal Diversity Web.
- FLORES-FOXWORTH, G., COONROD, S. A., MORENO, J. F., BYRD, S. R., KRAEMER, D. C. & WESTHUSIN, M. (1995). Interspecific transfer of IVM IVF-derived red sheep (*Ovis orientalis gmelini*) embryos to domestic sheep (*Ovis aries*). *Theriogenology* **44**, 681-690.
- FURLEY, C. W. (1986). Reproductive parameters of African gazelles: gestation, first fertile matings, first parturition and twinning. *African Journal of Ecology* **24**, 121-128.
- GAGNON, M. & CHEW, A. E. (2000). preferences in extant African bovidae. *Journal of Mammalogy* **81**, 490-511.
- GARDNER, A. L. (1971). Postpartum estrus in a Red brocket deer, *Mazama americana*, from Peru. *Journal of Mammalogy* **52**, 623-624.
- GAUTIER-HION, A., EMMONS, L. H. & DUBOST, G. (1980). A comparison of the diets of three major groups of primary consumers of Gabon (primates, squirrels and ruminants). *Oecologia* **45**, 182-189.
- GEIST, V. (1998). *Deer of the world. Their evolution, behaviour, and ecology*. Stackpole Books, Mechanicsburg, PA, USA.
- GIOCOMETTI, M. & RATTI, P. (1994). On the reproductive performance of free-ranging alpine ibex (*Capra i. ibex*) at Albris (Grisons, Switzerland). *Zeitschrift für Säugetierkunde* **59**, 174-180.
- GRAY, G. G. & SIMPSON, C. D. (1980). *Ammotragus lervia*. *Mammalian Species*, 1-7.

- GREEN, M. J. B. (1987). Diet composition and quality in Himalayan musk deer based on fecal analysis. *The Journal of Wildlife Management* **51**, 880-892.
- GRZIMEK, B. (1979/1980). *Grzimeks Tierleben Säugetiere 4*. DTV, Munich, Germany.
- GUINNESS, F. E., CLUTTON-BROCK, T. H. & ALBON, S. D. (1978). Factors affecting calf mortality in Red deer (*Cervus elaphus*). *Journal of Animal Ecology* **47**, 817-832.
- HAAGENRUD, H. & MARKGREN, G. (1974). The timing of estrus in Moose (*Alces alces* L.) in a district of Norway. In *International Congress of Game Biologists*, vol. 11, pp. 71-78, Stockholm.
- HABIBI, K. (1997). Group dynamics of the Nubian ibex (*Capra ibex nubiana*) in the Tuwayiq Canyons, Saudi Arabia. *Journal of Zoology* **241**, 791-801.
- HARDENBERG, A. (2000). Male Alpine chamois occupy territories at hotspots before the mating season. *Ethology* **106**, 617.
- HARRIS, R. B. & MILLER, D. J. (1995). Overlap in summer habitats and diets of Tibetan Plateau ungulates. *Mammalia* **59**, 197-212.
- HENTTONEN, H., STUBBE, M., MARAN, T. & A., T. (2008). *Alces alces*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.
- HOFMANN, R. R., STREICH, W. J., FICKEL, J., HUMMEL, J. & CLAUSS, M. (2008). Convergent evolution in feeding types: Salivary gland mass differences in wild ruminant species. *Journal of Morphology* **269**, 240-257.
- ILLIGE, D. (1951). An analysis of the reproductive pattern of Whitetail deer in South Texas. *Journal of Mammalogy* **32**, 411-421.
- IRBY, L. R. (1979). Reproduction in Mountain reedbuck (*Redunca fulvorufula*). *Mammalia* **43**, 191-214.
- IUCN SSC ANTELOPE SPECIALIST GROUP. (2008). *Oryx leucoryx*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4. .
- JACKSON, J. E. & GIULIETTI, J. (1988). The food habits of Pampas deer *Ozotoceros bezoarticus celer* in relation to its conservation in a relict natural grassland in Argentina. *Biological Conservation* **45**, 1-10.
- JASS, C. N. & MEAD, J. I. (2004). *Capricornis crispus*. *Mammalian Species*, 1-10.
- JONES, K. E., BIELBY, J., CARDILLO, M., FRITZ, S. A., O'DELL, J., ORME, C. D. L., SAFI, K., SECHREST, W., BOAKES, E. H., CARBONE, C., CONNOLLY, C., CUTTS, M. J., FOSTER, J. K., GRENYER, R., HABIB, M., PLASTER, C. A., PRICE, S. A., RIGBY, E. A., RIST, J., TEACHER, A., BININDA-EMONDS, O. R. P., GITTLEMAN, J. L., MACE, G. M., PURVIS, A. & MICHENER, W. K. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, 2648-2648.
- JORDAN, J. W. (1976). Natality of Black-tailed deer in McDonald State Forest, Oregon. *Northwest Science* **50**, 108.
- KEY, N. (2003). *Rusa alfredi*, pp. http://animaldiversity.ummz.umich.edu/site/accounts/information/Rusa_alfredi.html. Animal Diversity Web.
- KINGDON, J. (1982). *East African mammals. An atlas of evolution in Africa, Vol. III. Part C (bovids)*. Academic Press, London, Great Britain.
- KINGSWOOD, S. C. & BLANK, D. A. (1996). *Gazella subgutturosa*. *Mammalian Species* **518**, 1-10.
- KINGSWOOD, S. C. & KUMAMOTO, A. T. (1996). *Madoqua guentheri*. *Mammalian Species*, 1-10.
- KINGSWOOD, S. C. & KUMAMOTO, A. T. (1997). *Madoqua kirkii*. *Mammalian Species*, 1-10.
- KIRKWOOD, J. K. (2007). Management and formula intake of young hand-reared Chinese water deer *Hydropotes inermis* and their growth compared with mother-reared fawns. *International Zoo Yearbook* **27**, 308.

- KITCHEN, D. W. (1974). Social behavior and ecology of the Pronghorn. *Wildlife Monographs*, 3-96.
- KRANZ, K. R. & LUMPKIN, S. (1982). Notes on the Yellow-backed duiker in captivity with comments on its natural history. *International Zoo Yearbook* **22**, 232-240.
- KRAUSMAN, P. R. & CASEY, A. L. (2007). *Addax nasomaculatus*. *Mammalian Species* **807**, 1-4.
- LANGBEIN, J. (1998). Investigations on periparturient behaviour in free-ranging Mouflon sheep (*Ovis orientalis musimon*). *Journal of Zoology* **244**, 553.
- LANGBEIN, J., STREICH, J. & SCHEIBE, K. M. (1998). Characteristic activity patterns of female Mouflons (*Ovis orientalis musimon*) in the lambing period. *Applied Animal Behaviour Science* **58**, 281-292.
- LENT, P. C. (1966). Calving and related social behavior in the Barren-ground caribou. *Zeitschrift für Tierpsychologie* **23**, 701-756.
- LENT, P. C. (1988). *Ovibos moschatus*. *Mammalian Species*, 1-9.
- LESLIE, D. M. & SCHALLER, G. B. (2009). *Bos grunniens*. *Mammalian Species* **836**, 1-17.
- LESLIE, D. M. & SHARMA, K. (2009). *Tetracerus quadricornis*. *Mammalian Species*, 1-11.
- LEUTHOLD, W. (1967). Beobachtungen zum Jugendverhalten von Kob-Antilopen. *Zeitschrift für Säugetierkunde* **32**, 59-63.
- LEUTHOLD, W. (1977). *African ungulates*. Springer-Verlag, New York, USA.
- LEUTHOLD, W. & LEUTHOLD, B. M. (1975). Temporal patterns of reproduction in ungulates of Tsavo East National Park, Kenya. *African Journal of Ecology* **13**, 159-169.
- LINNELL, J. D. C. & ANDERSEN, R. (1998). Timing and synchrony of birth in a hider species, the Roe deer (*Capreolus capreolus*). *Journal of Zoology* **244**, 497-504.
- LOISON, A., GAILLARD, J.-M., PÉLABON, C. & YOCOZ, N. G. (1999). What factors shape sexual size dimorphism in ungulates? *Evolutionary Ecology Research* **1**, 611.
- LOUDON, A. S. I. (1988). Cycles of antler and testicular growth in an aseasonal tropical deer (*Axis axis*). *Journal of Reproduction and Fertility* **83**, 729.
- LUMPKIN, S. & KRANZ, K. R. (1984). *Cephalophus sylvicultor*. *Mammalian Species*, 1-7.
- LUNDRIGAN, B. & OAS, R. (2003). *Elaphodus cephalophus*. In *Animal Diversity Web*. Animal Diversity Web.
- LUTZ, H. (2001). *Tragulus napu*. Animal Diversity Web.
- MANSKI, D. A. (1991). Reproductive behavior of Addax antelope. *Applied Animal Behaviour Science* **29**, 39-66.
- MARUYAMA, N., SUGIMORI, F., TOTAKE, Y. & MIURA, S. (1975). The snorting voice of the sika deer in relation to its spacing distribution. *Journal of the Mammalogical Society of Japan* **6**, 155-162.
- MEAD, J. I. (1989). *Nemorhaedus goral*. *Mammalian Species*, 1-5.
- MENDELSSOHN, H. & YOM-TOV, Y. (1987). The mammals of Israel. In *Plants and animals of the Land of Israel: An illustrated encyclopedia*, vol. 7, pp. 1-295. Ministry of Defence Publishing House and Society for the Protection of Nature, Tel Aviv, Israel.
- MILLER, D. (2002). *Bubalus depressicornis*. Animal Diversity Web.
- MISHRA, H. R. & WEMMER, C. (1987). The comparative breeding ecology of four cervids in Royal Chitwan National Park. In *Biology and management of the Cervidae* (ed. C. M. Wemmer), pp. 259-271. Smithsonian Institution Press, Washington, D.C.
- MODHA, K. L. & ELTRINGHAM, S. K. (1976). Population ecology of the Uganda kob (*Adenota kob thomasi* (Neumann)) in relation to the territorial system in the Rwenzori National Park, Uganda. *Journal of Applied Ecology* **13**, 453-473.
- MOE, S. R., RUTINA, L. P. & DU TOIT, J. T. (2007). Trade-off between resource seasonality and predation risk explains reproductive chronology in impala. *Journal of Zoology* **273**, 237-243.

- MORROW, C. J. (1999). Reproductive seasonality in the female Scimitarâ horned oryx (*Oryx dammah*). *Animal Conservation* **2**, 261.
- MURRAY, M. G. (1982). The Rut of Impala: aspects of seasonal mating under tropical conditions. *Zeitschrift für Tierpsychologie* **59**, 319-337.
- NATIONAL RESEARCH COUNCIL. (1983). *Little-known Asian animal with a promising economic future*. National Academic Press, Washington, DC.
- NEAS, J. F. & HOFFMANN, R. S. (1987). *Budorcas taxicolor*. *Mammalian Species* **277**, 1-7.
- NORDIN, M. (1978). Voluntary food intake and digestion by the Lesser mouse-deer. *The Journal of Wildlife Management* **42**, 185-187.
- OCHIAI, K. (1999). Diet of the Japanese serow (*Capricornis crispus*) on the Shimokita Peninsula, northern Japan, in reference to variations with a 16-year interval. *Mammal Study* **24**, 91-102.
- OGREN, H. A. (1962). The Barbary sheep, *Ammotragus lervia* (Pallas), of the Canadian River Gorge, New Mexiko, University of Southern California.
- OLECH, W. (2008). *Bison bonasus*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4. .
- OWEN-SMITH, N. (1997). Distinctive features of the nutritional ecology of browsing versus grazing ruminants. *Zeitschrift für Säugetierkunde* **62**, 176-191.
- PADMALAL, U. K. G. K., TAKATSUKI, S. & JAYASEKARA, P. (2003). Food habits of Sambar *Cervus unicolor* at the Horton Plains National Park, Sri Lanka. *Ecological Research* **18**, 775-782-782.
- PAPPAS, L. A. (2002). *Taurotragus oryx*. *Mammalian Species*, 1-5.
- PEI, K., TABER, R. D., O'GARA, B. W. & WANG, Y. (1995). Breeding cycle of the Formosan Reeves muntjac (*Muntiacus reevesi micrurus*) in northern Taiwan, Republic of China. *Mammalia* **59**, 223-228.
- PENFOLD, L. M. (2005). Reproductive physiology and artificial insemination studies in wild and captive Gerenuk (*Litocranius walleri walleri*). *Reproduction, Fertility and Development* **17**, 707.
- PERELBERG, A., SALTZ, D., BAR-DAVID, S., DOLEV, A. & YOM-TOV, Y. (2003). Seasonal and circadian changes in the home ranges of reintroduced Persian fallow deer. *The Journal of Wildlife Management* **67**, 485-492.
- POLÁK, J. & FRYNTA, D. (2009). Sexual size dimorphism in domestic goats, sheep, and their wild relatives. *Biological Journal of the Linnaean Society* **98**, 872-883.
- POLLARD, S. (1999). *Pudu puda*. In *Animal Diversity Web*, vol. December 01, 2010.
- PRAKASH, I. (1960). Breeding of mammals in Rajasthan Desert, India. *Journal of Mammalogy* **41**, 386-389.
- PUJANINGSIH, R. I., SUTRISNO, C. I., SUPRIONDHO, Y., MALIK, A., DJUWANTOKO, PUDYATMOKO, S., AMIR, M. A. & ARYANTO, S. (2009). Diet composition of Anoa (*Buballus* sp.) studied using direct observation and dung analysis method in their habitat. *Journal of the Indonesian Tropical Animal Agriculture* **34**, 223-228.
- RABIEI, A. (2010). *Dama mesopotamica*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.
- RACHLOW, J. L. & BOWYER, R. T. (1991). Intraannual variation in timing and synchrony of parturition in Dall's sheep. *Journal of Mammalogy* **72**, 487-492.
- RALLS, K., KRANZ, K. & LUNDRIGAN, B. (1986). Mother-young relationships in captive ungulates: variability and clustering. *Animal Behaviour* **34**, 134-145.
- RALLS, K., LUNDRIGAN, B. & KRANZ, K. (1987). Mother-young relationships in captive ungulates: Spatial and temporal patterns. *Zoo Biology* **6**, 11-20.
- REAVILL, C. (2000). *Ovis aries*. Animal Diversity Web.

- REIMERS, E., KLEIN, D. R. & SØRUMGÅRD, R. (1983). Calving time, growth rate, and body size of Norwegian reindeer on different ranges. *Arctic and Alpine Research* **15**, 107-118.
- REYES, E. (2002). *Rusa timorensis*. Animal Diversity Web.
- RICE, C. G. (1988). Reproductive biology of Nilgiri tahr (*Hemitragus hylocrius*). *Journal of Zoology* **214**, 269-284.
- RICHTER. (1971). Observations on the biology and ecology of the Black wildebeest (*Connochaetes gnou*). *Journal of Southern African Wildlife Management Association* **1**, 3-16.
- RIDEOUT, C. B. & HOFFMANN, R. S. (1975). *Oreamnos americanus*. *Mammalian Species*, 1-6.
- ROBERTS, T. J. (1977). *The mammals of Pakistan*. Ernest Benn Limited, London, United Kingdom.
- RUTBERG, A. T. (1984). Birth synchrony in American bison (*Bison bison*): response to predation or season? *Journal of Mammalogy* **65**, 418-423.
- SANTIAGO-MORENO, J., LOPEZ-SEBASTIAN, A., GONZALEZ-BULNES, A., GOMEZ-BRUNET, A. & TORTONESE, D. (2001). The timing of the onset of puberty, extension of the breeding season, and length of postpartum anestrus in the female Mouflon (*Ovis gmelini musimon*). *Journal of Zoo and Wildlife Medicine* **32**, 230-235.
- SAYER, J. A. & LAVIEREN, L. P. V. (1975). The ecology of the Kafue lechwe population of Zambia before the operation of hydro-electric dams on the Kafue River. *African Journal of Ecology* **13**, 9-37.
- SCHALLER, G. B. (1967). The deer and the tiger. A study of wildlife in India. *University of Chicago Press Chicago*.
- SEKULIC, R. (1978). Seasonality of reproduction in the Sable antelope. *African Journal of Ecology* **16**, 177-182.
- SHACKLETON, D. M. (1985). *Ovis canadensis*. *Mammalian Species*, 1-9.
- SHENG, H. L. & LU, H. J. (1982). Distribution, habits and resource status of the Tufted deer (*Elaphodus cephalophus*). *Acta Zoologica Sinica* **28**, 307-311.
- SIMPSON, C. D. (1968). Reproduction and population structure in Greater kudu in Rhodesia. *The Journal of Wildlife Management* **32**, 149-161.
- SINCLAIR, A. R. E., MDUMA, S. A. & ARCESE, P. (2000). What determines phenology and synchrony of ungulate breeding in Serengeti? *Ecology* **81**, 2100-2111.
- SINGH, G., SINGH, G. B. & BAINS, G. S. (1985). Studies on incidence of anoestrus in village buffaloes [India]. *Indian Journal of Animal Research* **19**, 57-60.
- SKINNER, J. D., AARDE, R. J. V., KNIGHT, M. H. & DOTT, H. M. (1996). Morphometries and reproduction in a population of Springbok *Antidorcas marsupialis* in the semi-arid southern Kalahari. *African Journal of Ecology* **34**, 312-330.
- SKINNER, J. D., MOSS, D. G. & SKINNER, D. C. (2002). Inherent seasonality in the breeding seasons of African mammals: evidence from captive breeding. *Transactions of the Royal Society of South Africa* **57**, 25.
- SOKOLOV, V. E. (1974). *Saiga tatarica*. *Mammalian Species*, 1-4.
- SOLANKI, G. S. & NAIK, R. M. (1998). Grazing interactions between wild and domestic herbivores. *Small Ruminant Research* **27**, 231-235.
- SPINAGE, C. A. (1973). The role of photoperiodism in the seasonal breeding of tropical African ungulates. *Mammal Review* **3**, 71-83.
- SPINAGE, C. A. (1986). *The natural history of antelopes*. Croom Helm Publishers Ltd, London.
- SPONHEIMER, M., LEE-THORP, J. A., DERUITER, D. J., SMITH, J. M., VAN DER MERWE, J., GRANT, C. C., AYLIFFE, L. K., ROBINSON, T. F., HEIDELBERGER, C. & MARCUS, W. (2003). Diets of southern African Bovidae: stable isotope evidence. *Journal of Mammalogy* **84**, 471.

- STERBA, O. & KLUSAK, K. (1984). Reproductive biology of Fallow deer, *Dama dama*. *Acta Scientifica Nature Borneo* **18**, 1-46.
- STONER, C. J., CARO, T. M. & GRAHAM, C. M. (2003). Ecological and behavioral correlates of coloration in artiodactyls: systematic analyses of conventional hypotheses. *Behavioral Ecology* **14**, 823-840.
- STRINGHAM, S. F. (1974). Mother-infant relations in semi-captive Alaskan mosse (*Alces alces gigas*), University of Alaska.
- STÜWE, M. & GRODINSKY, C. (1987). Reproductive biology of captive Alpine ibex (*Capra i. ibex*). *Zoo Biology* **6**, 331-339.
- SUMITHRAN, S. (1997). Status and ecology of the Nilgiri tahr in the Mukurthi National Park, South India. In Fisheries and Wildlife Sciences, Virginia Polytechnic Institute and State University.
- TOM-TOV, Y. (1995). *Gazella dorcas*. *Mammalian Species* **491**, 1-6.
- TURKALO, A. & TURKALO. (1999). Group size and group composition of the Bongo (*Tragelaphus eurycerus*) at a natural lick in the Dzanga National Park, Central African Republic. *Mammalia* **63**, 437.
- UNGERFELD, R., GONZALEZ-PENSADO, S., BIELLI, A., VILLAGRAN, M., OLAZABAL, D. & PEREZ, W. (2008). Reproductive biology of the Pampas deer (*Ozotoceros bezoarticus*): a review. *Acta Veterinaria Scandinavica* **50**, 16.
- UNKNOWN AUTHOR. (2005). The hoofed mammals. In *The Endangered Mammals of Tibet*, pp. 1-40. DIIR.
- VAIRAVEL, S. M. (1998). Ecology of Gaur (*Bos gaurus*) with special reference to habitat utilization in Parambikulam Wildlife Sanctuary, Kerala, India.
- VALDEZ, R. (2008). *Ovis orientalis*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.
- VALDEZ, R., CARDENAS, M. & SANCHEZ, J. (1991). Disruptive mating behavior by subadult Armenian wild sheep in Iran. *Applied Animal Behaviour Science* **29**, 165-171.
- VAN WIEREN, S. E. (1996). Browsers and grazers: foraging strategies in ruminants. In *Digestive strategies in ruminants and nonruminants* (ed. S. E. Van Wieren), pp. 119-146. Thesis Landbouw Universiteit Wageningen, Wageningen, The Netherlands.
- VIÉ, J. C. (1996). Reproductive biology of captive Arabian oryx (*Oryx leucoryx*) in Saudi Arabia. *Zoo Biology* **15**, 371.
- VRAHIMIS, S. & KOK, O. B. (1994). Notes on the diurnal activity of early post-natal black wildebeest calves. *Koedoe* **37**, 109-113.
- WANG, X. & HOFFMANN, R. S. (1987). *Pseudois nayaur* and *Pseudois schaeferi*. *Mammalian Species*, 1-6.
- WANG, X. Z., SONG, Y. L., ZENG, Z. G., GONG, H. S. & ZHAO, N. Z. (2005). Characteristics of birth-site and early mother-infant relationship of Golden takin *Budorcas taxicolor bedfordi*. *Acta Zoologica Sinica* **51**, 748-752.
- WATSON, L. H. & OWEN-SMITH, N. (2002). Phenological influences on the utilization of woody plants by eland in semi-arid shrubland. *African Journal of Ecology* **40**, 65-75.
- WECKERLY, F. W. (1998). Sexual-size dimorphism: influence of mass and mating systems in the most dimorphic mammals. *Journal of Mammalogy* **79**, 33-52.
- WEGGE, P., SHRESTHA, A. K. & MOE, S. R. (2006). Dry season diets of sympatric ungulates in lowland Nepal: competition and facilitation in alluvial tall grasslands. *Ecological Research* **21**, 698-706.
- WEINBERG, P. J. (2002). *Capra cylindricornis*. *Mammalian Species* **295**, 1-9.
- WEINBERG, P. J. (2004). West Caucasian Tur (*Capra [ibex] caucasica* Güld. et all.), biology, status and taxonomy. In *2nd International Conference on Alpine ibex*, Cogne, Italy.
- WEINBERG, P. J. (2008). *Capra cylindricornis*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.

- WEMMER, C., HALVERSON, T., RODDEN, M. & PORTILLO, T. (1989). The reproductive biology of female Père David's deer (*Elaphurus davidianus*). *Zoo Biology* **8**, 49-55.
- WILSON, D. E. & HIRST, S. M. (1977). Ecology and factors limiting Roan and Sable antelope populations in South Africa. *Wildlife Monographs*, 3-111.
- WILSON, V. J. (2005). *Duikers of Africa (Masters of the African forest floor)*. Zimbi Books, Pretoria (South Africa).
- WORREL, E. (2004). *Rucervus eldii*. Animal Diversity Web.
- YOM-TOV, Y., MENDELSSOHN, H. & GROVES, C. P. (1995). *Gazella dorcas*. *Mammalian Species*, 1-6.
- ZHIGANG, J. & HARRIS, R. B. (2008). *Elaphurus davidianus*. IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4.
- ZI, X.-D. (2003). Reproduction in female yaks (*Bos grunniens*) and opportunities for improvement. *Theriogenology* **59**, 1303-1312.

Curriculum vitae

Name	Philipp Zerbe
Geburtsdatum	07.05.1985
Geburtsort	Offenbach am Main
Nationalität	Deutsch
1992-1996	Grundschule (Kolpingschule, Aschaffenburg, Deutschland)
1996-2002	Friedrich-Dessauer-Gymnasium (Aschaffenburg, Deutschland)
2002-2004	Kantonsschule Reussbühl (Kanton Luzern, Schweiz)
17.06.2004	Matura, Kantonsschule Reussbühl (Kanton Luzern, Schweiz) Schwerpunkt: Mathematik-Physik
2004-2010	Studium Veterinärmedizin, Vetsuisse-Fakultät Zürich, Schweiz
14.10.2010	Diplom Veterinärmedizin, Vetsuisse-Fakultät Zürich, Schweiz
2010-2012	Anfertigung der Dissertation unter Leitung von PD Dr. med. vet. Marcus Clauss am Departement für Kleintiere, Klinik für Zoo-, Heim- und Wildtiere Vetsuisse-Fakultät Universität Zürich Direktor: Prof. Dr. med. vet. Jean-Michel Hatt
2011-2012	Internship/Assistenz Abteilung für Kleintierfortpflanzung Klinik für Fortpflanzungsmedizin, Universität Zürich, Schweiz
2012-	Veterinary Research Fellowship, Preclinical Service Focus Area Surgery, AO Research Institute Davos, Schweiz

Von tiefstem Herzen danke ich...

meinen Eltern Dagmar und Manfred für Eure stetige Unterstützung und Liebe, die Wurzeln und Flügel und ein zu Hause welches tief im Herzen sitzt.

meinen Brüdern Adrian und Simon für all die Herausforderungen, Erlebnisse und Kämpfe die wir zusammen durchgestanden haben. Danke für unser tiefes Vertrauen!

meinen Grosseltern Erika, Gitta und Roland für Eure Frühförderung im aktiven, sozialen und gesellschaftlichen Sinne. Danke dass Ihr immer da seid!

meinem Betreuer Marcus Clauss für die Motivation einen kleinen Studenten der eines Tages buchstäblich vor Deiner Türe stand in das Leben der Wissenschaft einzuführen und Deine Ausdauer und Vertrauen, die Du dabei an den Tag gelegt hast.

meinem Betreuer Dennis Müller für Deine aufbauenden Worte und Unterstützung bei jeder meiner noch so kleinen Ideen, die uns bis in die Nacht hinein beschäftigt haben.

meinem Mentor Karl Nuss für die Verbindung die irgendwie immer da war und mit Gesprächen ganze Abende gefüllt hat. Danke für die Zeit und Unterstützung zum genau richtigen Zeitpunkt.

meinem stillen Mentor Thomas Lutz für Deine aufbauenden Worte, welche immer wieder gebraucht wurden.

meinem Freund und Bruder Matthias „Mäd“ für unsere tiefe Verbundenheit.

meinem Freund und Bruder Yves für all die Jahre die Du neben mir in der WG ausgehalten und meinem Fahrrad das Fliegen beigebracht hast. Danke für jedes Gespräch zwischen Tür und Angel, in Kermit, den Lernpausen und überall.

meinem besten Freund Jan für die regelmässig jährlichen Treffen, welche uns jedes Mal in die alte Zeit zurückversetzt hat und sämtliche Erziehung hat vergessen lassen.

meiner namibianischen Familie Nandi, Kim, Jimmy, Ulrike und Helmut für eine unvergessliche Zeit, die mich auf Ewig geprägt hat und Euch immer einen Platz in meinem Herzen und damit auch meinem zu Hause sichert!

meiner WG 3F – Michi, Nora, Desi, Dani, Aurelien und allen anderen - für all das Theater, was mich immer abgelenkt hat. Ihr wart immer eine Family...

meinen guten Freunden Sylvia und Niggi die mich immer mit Herzen aufgenommen haben und ähnliche abstruse Ideen in die Tat umsetzen für all die unvergesslichen Momente die wir jetzt schon zusammen hatten. Und für

Chukela, die es die ganze Zeit unter meinem Schreibtisch im Doktorandenzimmer ausgehalten hat und auch nachts um 2 noch begeistert vom Büro nach Hause gelaufen ist.